

ISOMERIC METASTABLE STATES IN
MEDIUM AND HEAVY WEIGHT NUCLEI

—♦♦♦—
A. E. FRANCIS

1951

Library
U. S. Naval Postgraduate School
Monterey, California

ISOMERIC METASTABLE STATES IN
MEDIUM AND HEAVY WEIGHT NUCLEI

by

Author Eugene Francis

M. E. Stevens Institute of Technology
(1942)

M. S. Massachusetts Institute of Technology
(1948)

Submitted in Partial Fulfillment of the
Requirements for the Degree of
Master of Science in Physics

at the

Massachusetts Institute of Technology
(1951)

Signature of Author _____
Department of Physics, July 1, 1951

Certified by _____
Thesis Supervisor

Chairman, Departmental Committee on Graduate Students

THE UNITED STATES OF AMERICA
DEPARTMENT OF AGRICULTURE

19

UNITED STATES DEPARTMENT OF AGRICULTURE
BUREAU OF PLANT INDUSTRY
(1917)
UNITED STATES DEPARTMENT OF AGRICULTURE
(1917)

UNITED STATES DEPARTMENT OF AGRICULTURE
BUREAU OF PLANT INDUSTRY
UNITED STATES DEPARTMENT OF AGRICULTURE
BUREAU OF PLANT INDUSTRY

UNITED STATES DEPARTMENT OF AGRICULTURE
(1917)

UNITED STATES DEPARTMENT OF AGRICULTURE
BUREAU OF PLANT INDUSTRY

UNITED STATES DEPARTMENT OF AGRICULTURE
BUREAU OF PLANT INDUSTRY

UNITED STATES DEPARTMENT OF AGRICULTURE
BUREAU OF PLANT INDUSTRY

ISOMERIC METASTABLE STATES IN MEDIUM AND HEAVY WEIGHT NUCLEI

by

Arthur Eugene Francis

Submitted for the Degree of Master of Science in the Department
of Physics on July 17, 1951

ABSTRACT

The 48.6-minute metastable state was formed in Cd^{111} by the inelastic scattering of monoenergetic fast neutrons on natural cadmium. The neutrons were obtained by bombarding a Li^7 target with monoenergetic protons from the Rockefeller generator at M.I.T. The presence of excited levels in Cd^{111} at energies of 0.720 ± 0.020 Mev and 1.150 ± 0.020 Mev was detected. The threshold for the formation of the metastable state was determined as 0.400 ± 0.25 Mev. Therefore, either the 0.420-Mev level or the 0.396-Mev, 48.6-minute, metastable level was excited directly. Either an $l = 3$ or an $l = 4$ neutron combines with Cd^{111} to form the compound nucleus which emits an $l = 0$ neutron. If the 48.6-minute metastable state was excited directly, and that is almost certain, its angular momentum is either $11/2$ or $9/2$; the angular momentum of the 5×10^{-8} -second metastable state is either $5/2$ or $3/2$; the parities of the two metastable states are opposite to the parity of the ground state if the incoming neutron has $l = 3$; and the parities are the same as the parity of the ground state if the incoming neutron has $l = 4$.

THE UNITED STATES OF AMERICA

OF

THE DISTRICT OF COLUMBIA

IN SENATE, JANUARY 11, 1901

REPORT

OF

THE COMMISSIONERS OF THE DISTRICT OF COLUMBIA

IN RESPONSE TO A RESOLUTION PASSED BY THE SENATE

ON JANUARY 11, 1901

AND PRINTED BY THE GOVERNMENT PRINTING OFFICE

WASHINGTON: 1901

THE DISTRICT OF COLUMBIA

IN SENATE, JANUARY 11, 1901

REPORT

OF

THE COMMISSIONERS OF THE DISTRICT OF COLUMBIA

IN RESPONSE TO A RESOLUTION PASSED BY THE SENATE

ON JANUARY 11, 1901

AND PRINTED BY THE GOVERNMENT PRINTING OFFICE

WASHINGTON: 1901

THE DISTRICT OF COLUMBIA

IN SENATE, JANUARY 11, 1901

2

TABLE OF CONTENTS

	Page
ABSTRACT	i
LIST OF FIGURES	iv
LIST OF TABLES	vi
CHAPTER I. INTRODUCTION	1
CHAPTER II. THEORY	6
Half-Lives of Excited States	6
Selection Rules for Gamma-Ray Transitions	6
Internal Conversion	8
Neutron Excitation of Metastable Levels	10
Neutron Versus Photon Excitation of Metastable States	14
CHAPTER III. EXPERIMENTAL EQUIPMENT	16
Source of Neutrons	16
Rockefeller Generator	17
Measurement of Neutron Yield	18
Scintillation Counter	21
Platinum-Screen Counter	22
CHAPTER IV. INVESTIGATION OF Cd ¹¹¹	23
Historical Background of the Excitation of Metastable Cd ¹¹¹	23
Energy Levels in Cd ¹¹¹	24
Method of Obtaining the Uncorrected Excitation Curve	31
Elimination of the Effect of Interfering Decays	40

TABLE OF CONTENTS

1	INTRODUCTION
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	
25	
26	
27	
28	
29	
30	
31	
32	
33	
34	
35	
36	
37	
38	
39	
40	
41	
42	
43	
44	
45	
46	
47	
48	
49	
50	
51	
52	
53	
54	
55	
56	
57	
58	
59	
60	
61	
62	
63	
64	
65	
66	
67	
68	
69	
70	
71	
72	
73	
74	
75	
76	
77	
78	
79	
80	
81	
82	
83	
84	
85	
86	
87	
88	
89	
90	
91	
92	
93	
94	
95	
96	
97	
98	
99	
100	

	Page
Correction of the 45.6-Minute Excitation Curve for Proton Energy Spread, Target Thickness, and Angularity	47
Determination of the Absolute Cross Section for the Formation of the Metastable State	52
Comparison of the Theoretical and Experimental Cross Sections	60
CHAPTER V. CONCLUSIONS AND SUGGESTIONS FOR FURTHER INVESTIGATION	66
Discussion and Summary of Results	66
Suggestions for Further Investigation	69
BIBLIOGRAPHY	72
ACKNOWLEDGMENTS	75
APPENDIX A. ALTERNATION OF THE SINGLE CHANNEL DIFFERENTIAL DISCRIMINATOR	76
APPENDIX B. DETERMINATION OF THE ABSOLUTE RATE OF GAMMA RAY EMISSION OF THE ^{203}Hg STANDARD	77
APPENDIX C. UNSUCCESSFUL PRELIMINARY EXPERIMENTS IN PRODUCING AND DETECTING THE DECAY OF META- STABLE STATES	80

1. The first of these is the fact that the Commission has not yet received any information from the Government of the United States regarding the activities of the Committee for the Liberation of the People of the South (CLPS) in the United States.

2. The second is the fact that the Commission has not yet received any information from the Government of the United States regarding the activities of the Committee for the Liberation of the People of the South (CLPS) in the United States.

3. The third is the fact that the Commission has not yet received any information from the Government of the United States regarding the activities of the Committee for the Liberation of the People of the South (CLPS) in the United States.

4. The fourth is the fact that the Commission has not yet received any information from the Government of the United States regarding the activities of the Committee for the Liberation of the People of the South (CLPS) in the United States.

5. The fifth is the fact that the Commission has not yet received any information from the Government of the United States regarding the activities of the Committee for the Liberation of the People of the South (CLPS) in the United States.

6. The sixth is the fact that the Commission has not yet received any information from the Government of the United States regarding the activities of the Committee for the Liberation of the People of the South (CLPS) in the United States.

7. The seventh is the fact that the Commission has not yet received any information from the Government of the United States regarding the activities of the Committee for the Liberation of the People of the South (CLPS) in the United States.

8. The eighth is the fact that the Commission has not yet received any information from the Government of the United States regarding the activities of the Committee for the Liberation of the People of the South (CLPS) in the United States.

9. The ninth is the fact that the Commission has not yet received any information from the Government of the United States regarding the activities of the Committee for the Liberation of the People of the South (CLPS) in the United States.

10. The tenth is the fact that the Commission has not yet received any information from the Government of the United States regarding the activities of the Committee for the Liberation of the People of the South (CLPS) in the United States.

LIST OF FIGURES

		Page
Figure 1.	Schematic Diagram of Neutron Scattering	11
Figure 2.	Long Counter	19
Figure 3.	Response of Long Counter to Monoenergetic Neutrons	20
Figure 4.	Energy Level Diagram for Cd^{111}	25
Figure 5.	Possible Values of Parity and Angular Momentum for the Metastable States of Cd^{111}	30
Figure 6.	Arrangement for Irradiating the Cadmium Disk, Showing the Magnet and Chamber of the Rockefeller Generator, the Tantalum-Backed Lithium Target, and the Aluminum Disk Holder	35
Figure 7.	Setup of Equipment Used for Detecting the Decay of the Metastable State, Showing the High-Voltage Power Supply, Voltage Stabilizer, Lead-Sheathed Scintillation Counter, Cathode-Follower Pre-amplifier, Amplifier, and the Decade Scaler	36
Figure 8.	Neutron Excitation Curve for ^{111}Cd - Uncorrected	39
Figure 9.	Basic Decay Curve for Irradiation by 0.532-Mev Neutrons	42
Figure 10.	Comparison of Decays for Different Percentages of 48.6-Minute Activity	44
Figure 11.	Excitation Curves for the Separated Activities ^{111}Cd , ^{115}Cd , and ^{117}Cd	46

LIST OF FIGURES

Figure 1.	Automatic system of control	11
Figure 2.	Control system	12
Figure 3.	Diagram of test system in laboratory	13
Figure 4.	Block diagram of test system	14
Figure 5.	Block diagram of test system and system for the automatic control of the test system	15
Figure 6.	Diagram of test system and system for the automatic control of the test system	16
Figure 7.	Diagram of test system and system for the automatic control of the test system	17
Figure 8.	Diagram of test system and system for the automatic control of the test system	18
Figure 9.	Diagram of test system and system for the automatic control of the test system	19
Figure 10.	Diagram of test system and system for the automatic control of the test system	20
Figure 11.	Diagram of test system and system for the automatic control of the test system	21
Figure 12.	Diagram of test system and system for the automatic control of the test system	22

Figure 12.	Excitation Curve for 45.6-Minute Metastable ^{111}Cd Corrected for Finite Target Thickness and Angularity of the Incident Neutrons	53
Figure 13.	Cross Section for the Formation of Metastable State in ^{111}Cd	61
Figure 14.	Comparison of the Experimental Cross Section for Formation of the Metastable State with the Theoretical Cross Section for Formation of the Compound Nucleus	63
Figure 15.	Probability of Compound Nucleus Decaying to the Metastable State	65
Figure 16.	Most Probable Assignment of Angular Momentum and Parity to the Metastable States of ^{111}Cd .	70
Figure 17.	Calibrated Curve for Standardising the Hg^{203} Sample	78

LIST OF TABLES

	Page
TABLE I. The Theoretically Calculated Half-Lives for Decay by Gamma-Ray Emission of Different Multipole Orders for the Metastable States of Cd ¹¹¹	26
TABLE II. Data on Internal Conversion of Cd ¹¹¹	28
TABLE III. Isotopes of Cadmium	33

I. INTRODUCTION

The determination of the energy, angular momentum, and parity of excited levels in nuclei is an important step in obtaining information concerning nuclear structure.

These characteristics of excited levels can be obtained from observations of transitions between specific excited states and between excited states and the ground state in nuclei. Such transitions often occur after a nuclear reaction leaves the nucleus in an excited state. However, these excited states usually have very short lifetimes.

Nuclear isomers are nuclei having the same number of protons and neutrons, but differing in their stability or mode of radioactive decay. The simplest case of nuclear isomerism occurs when one of the isomers is a stable nucleus in its ground state and the other is an excited state of the same nucleus. When an excited state exists for a measurable length of time, it is called a metastable state. Metastable states differ from other excited states only because of their slower rate of decay.

In 1921, Otto Hahn, the German radiochemist, produced evidence that uranium Z and uranium X_2 , both naturally occurring nuclides, were isotopic and isobaric, but had different radioactive decay rates. For several years, this remained as the only substantiated case of nuclear isomerism. In 1935, it was discovered that the bombardment of $_{35}\text{Br}^{79}$ by slow neutrons produced two isomers of Br^{80} with different half-lives. Since that time, many examples of nuclear isomerism have

been discovered. As measurement techniques improve, more radioactive nuclides are being reclassified as isomeric.

Weizsacker provided the first theoretical explanation for the existence of isomers. He showed that by assigning to the excited state an angular momentum which differs from the ground state by 3 to 5 units and an energy of a few hundred kilovolts or less, a half-life is predicted for transition to the ground state by gamma-ray emission which is long enough to be observable.

Isomeric metastable states result from interactions between high-energy X-rays or electrons and stable nuclei and from the nuclear excitation which accompanies the inelastic scattering of alpha particles, deuterons, protons, or fast neutrons. Inelastic scattering of neutrons is a convenient method of producing excited states in nuclei, but cross sections for such reactions are small, and large inherent backgrounds are encountered. However, if the lifetime of the excited state is long enough, the transition to a lower state may be observed after the beam of exciting particles has been removed and after competing reactions have died away. Therefore, the existence of metastable states provides a method of studying energy levels in nuclei. Comparison of the experimentally obtained cross section for the formation of the metastable state in a particular nucleus with the theoretical cross section for the formation of the same state often facilitates estimating the spin and parity of the metastable state and of other excited levels in that nucleus.

weibungsbereit sein, wenn ein Individuum i die Entscheidung trifft

Copyright © 1999 by John Wiley & Sons, Inc.

Yell. gelatinöse, bei 10°C wasserig, bei 20°C dickflüssig.

Please do not write on this form.

10-10-68

© 2000 Blackwell Science Ltd, *Journal of Internal Medicine* 247: 395–402

to provide the following information:

While it has been noted that the

Immunization with *Yersinia enterocolitica* 103

This document will expire June 30, 2006. See [http://www.fda.gov/cder/rtx/rtx.htm](#) for more information.

...the

...to the

Source: Inf. Values of private business activities. In: *Business & Economics* 4, 1977, p. 10.

Abstracted from: *Journal of the American Academy of Child and Adolescent Psychiatry*, 1995, 34(10), 1241-1250.

...and the ...

Source: The author's calculations based on data from the 1990 Census.

willkommen willkommen heißt den Besucher und den Besucher willkommen. Es

— the fact that the system is not a simple one, and that it is not a simple one.

© 2000 Blackwell Science Ltd *Journal of Internal Medicine* 247: 395–401

1987-1988 season. The following table shows the results of the study.

the following - the same type of information for the

THIS IS NOT A RACIAL MATTER. IT IS A MATTER OF HUMAN RIGHTS.

The half-lives of the various known isomers range from fractions of seconds to years, and there is no qualitative difference between an excited state with a very short life and one with a long life. Isomers have not been found in nuclei having mass numbers below 39. The characteristics (half-life, excitation energy, decay scheme, conversion ratios, etc.) of isomers of stable nuclei may be found in various tabulations of nuclear data, such as Seaborg and Pearlman (S1), Sullivan (S2), and Nuclear Data NBS 499 (N1). Segre and Helmholtz (S3) have assembled complete data on all but the most recently discovered isomers. A careful survey of their tables is required when selecting a particular isomer for experimental investigation. The experimenter must be sure that there are no other radiations arising from accompanying nuclear reactions which seriously interfere with the detection of the decay of the metastable state.

When the activity of a metastable state is plotted against the energy of the exciting medium, changes in the slope of the curve occur as new excited levels of the stable nucleus are reached. The excitation curve changes shape because additional higher excited levels in the target nucleus become available for decay to the metastable state as the excitation energy increases, and an increase in the cross section for formation of the metastable state is produced. Wiedenbeck (W1, W2, W3, W4) has obtained such excitation curves which indicate excited levels that decay to the metastable state for Au¹⁹⁷, Rh¹⁰³, Cd¹¹¹, and both isotopes of Ag. Wiedenbeck (W4) also reports excitation of

the following is a list of the names of the persons who have been appointed to the various committees of the National Council on the Status of Women, established by the Executive Order of the President on June 1, 1946.

the threshold level alone in Au^{197} by bombardment with fast neutrons of broad energy distribution, and he states that the neutron threshold is the same as the one obtained in his X-ray work.

Waldman and Wiedenbeck (45), using X-rays as an exciting medium, have reported a series of excited levels which decay to the metastable state in In^{115} . Goldhaber, Hill, and Szilard (61) have confirmed the existence of a metastable state in In^{115} by inelastic scattering of fast neutrons. The metastable state in In^{115} has also been excited by Barnes and Aradine (51) with protons and by Lark-Horovitz, Riasser, and Smith (11) with high-energy alpha particles. Both Cohen (61) and Taschek (71) have obtained excitation curves for In^{115} , bombarded by fast neutrons which indicate only the threshold level.

The quanta employed in the foregoing work involving the use of X-rays were obtained by bombarding thick targets with monoenergetic electrons. Consequently, the X-rays were continuous in energy distribution. Prior to 1950, no experimenters had used monoenergetic heavy particles to determine excited levels by inelastic scattering for an appreciable range of energies above the metastable state. The use of either quanta or heavy particles with a substantial energy distribution results in only slight changes in the slope of the excitation curve at energy levels. Therefore, the exact location and sometimes even the existence of excited levels is subject to question. Abel (51), using monoenergetic fast neutrons

as a source of excitation, located several energy levels in Au¹⁹⁷ and In¹¹⁵. The changes in slope of the monoenergetic neutron excitation curves at energy levels are pronounced enough to inspire considerable confidence. All the levels determined by Ebel appear to be slightly lower than the corresponding levels previously determined by quantum excitation. This thesis describes the use of Ebel's technique in investigating the metastable state and higher energy excited states in Cd¹¹¹.

II. THEORY

Half-Lives of Excited States

Segre and Helmholtz (53) have presented a detailed review of Weissacker's theory of nuclear isomerism. Another good survey of various theories of nuclear isomerism is given by Hole (H1). Such theoretical quantitative explanations for the measurable lifetimes of metastable states are complicated by the dearth of detailed information about the nucleus. Consequently, many different methods for calculating half-lives of metastable states have been suggested, each justified by different assumptions concerning the nature of the nucleus. Bethé (B2), Hebb and Uhlenbeck (E2), Lowen (L2), and others have determined formulas for predicting the probability of transition from an excited level to a lower level by gamma-ray emission. Moon (M1) gives a graph, based on a simplification of Lowen's formula, of decay constants vs. gamma-ray energy for various multipole values of radiation. This curve is a convenient method of estimating either the half-life, the multipole order, or the transition energy of a particular gamma-ray transition when two of three parameters are known quantities.

Selection Rules for Gamma-Ray Transitions

The general theory regarding gamma-ray emission (selection rules, decay constants, spin, parity, etc.) includes transitions from

the metastable state. The quantum theory of radiation utilizes the classical concept of a radiation source as an oscillating electric or magnetic moment. The radiations are classified as 2^l electric or magnetic dipole, quadrupole, octupole, etc., where l is the angular momentum carried away by the gamma ray in units of \hbar . Since the vector angular momentum of the system is conserved in all nuclear reactions, the emission of a gamma ray by a nucleus changes its angular momentum in accordance with the multipole order of the radiation. The absolute difference in angular momentum between two states in a nucleus determines the lowest allowed multipole order radiation that can occur in a transition between the states ($l_{\min} = |I - I'|$). Higher order multipole radiations up to $l_{\max} = |I + I'|$ are possible. However, the probability per unit time of radiation occurring decreases rapidly as the multipole order increases, so the lowest allowed multipole order is usually the only one which contributes appreciably to the transition, except that an electric 2^{l+1} -pole may be a serious competitor to a magnetic 2^l -pole.

The parity of a state is said to be odd (-) or even (+) depending on whether a reversal of sign of all the coordinates of the system does or does not reverse the sign of the wave function describing the state. Parity, like angular momentum, is conserved in a nuclear reaction. The parity of even electric and odd magnetic multipoles is taken as even (+) and the parity of odd electric and even magnetic multipoles is taken as odd (-). Therefore, if two states have the same

The following table shows the results of the analysis of the data obtained from the experiments. The results are given in the form of a table, the columns of which are headed by the names of the variables, and the rows by the names of the experiments. The results are given in the form of a table, the columns of which are headed by the names of the variables, and the rows by the names of the experiments.

parity, a transition between them involves only even electric or odd magnetic multipole orders, and if two states have opposite parity, the transition between them involves only odd electric and even magnetic multipole orders.

The importance of the angular momentum and parity selection rules is manifested in the long lifetime of a metastable state.

Internal Conversion

A nucleus in an excited state may spontaneously change to a state of lower energy by emitting a gamma ray of energy equal to the energy difference between the two states or by giving this increment of energy to an atomic electron in one of its own shells. This "internal conversion" electron is ejected from the atom with a kinetic energy equal to the corresponding quantum energy less the atomic binding energy of the electron and the energy of recoil. Originally it was thought that the energy of the nuclear transition was first emitted as a gamma ray, which then ejected a photo-electron from the atom. Taylor and Kott (21) showed that internal conversion is actually a primary rather than secondary process and is due to the direct interaction of the nucleus with its own electrons. Therefore, the probability per unit time of decay of an excited state may be the sum of the probabilities of the competing gamma-ray emission and internal conversion instead of the probability of gamma-ray emission only. The probability of internal conversion may be even greater than that

1. The first of these is the fact that the
2. second is the fact that the
3. third is the fact that the
4. fourth is the fact that the
5. fifth is the fact that the
6. sixth is the fact that the
7. seventh is the fact that the
8. eighth is the fact that the
9. ninth is the fact that the
10. tenth is the fact that the

1. The following is a list of the names of the persons who have been appointed to the various committees of the Board of Directors of the American Telephone and Telegraph Company, for the year ending December 31, 1910.

It is a very common mistake to think that the only way to get a good result is to use a lot of force. In fact, the best results are often achieved by using a small amount of force, applied in a precise and controlled manner. This is why it is so important to have a good understanding of the mechanics of the system you are working with. Only by knowing exactly how the system works can you apply the right amount of force at the right time and in the right direction. This is the key to success in any technical field.

1. The first step in the process of identifying a problem is to define the problem. This involves identifying the symptoms of the problem and determining the scope of the problem. Once the problem has been defined, the next step is to identify the causes of the problem. This involves identifying the factors that are contributing to the problem and determining the underlying causes. Once the causes have been identified, the next step is to develop a plan of action. This involves identifying the steps that need to be taken to solve the problem and determining the resources that will be needed to implement the plan. Once a plan of action has been developed, the final step is to implement the plan. This involves carrying out the steps that have been identified in the plan and monitoring the progress of the implementation. Once the problem has been solved, the final step is to evaluate the results of the implementation. This involves determining whether the problem has been solved and whether the resources have been used effectively.

of gamma-ray emission. The ejection of the conversion electron causes a vacancy in the electronic structure of the atom, and therefore, it is followed immediately by the emission of characteristic X-rays of the element in question.

The conversion coefficient is the ratio of the rate of emission of conversion electrons to the rate of emission of gamma rays. The total rate of electron emission is the sum of the rates for the various atomic shells (K, L, etc.). The conversion coefficient is complicated to calculate theoretically and difficult to determine experimentally. Dancoff and Morrison (D1) have obtained an expression for K-electron conversion coefficients which treats the ejected electron nonrelativistically and have also obtained an approximate relativistic expression which neglects electron binding energy. Another source of K-shell internal conversion coefficients is an extensive table which has been privately distributed by Rose (R1). The ratio of the K-shell to the L-shell conversion coefficients is relatively easy to determine experimentally because it is only necessary to compare the intensity of two lines in the same electron spectrum. Comparison of experimental and theoretical K/L ratios affords a means of determining whether a gamma-ray transition is an electric or magnetic multipole.

Axel and Dancoff (A1) have found good agreement between the experimentally determined lifetimes of about 50 isomers and the theoretical lifetimes predicted by the Weissacker hypothesis and corrected for internal conversion.

1. The first step is to identify the problem or question that needs to be answered. This involves understanding the context and the specific requirements of the task.

[illegible]

Neutron Excitation of Metastable Levels

Excitation of a metastable state in a nucleus by bombarding the nucleus with fast neutrons is a special case of inelastic scattering of fast neutrons. The process is shown diagrammatically in Figure 1. Bombardment by neutrons with less energy than the energy of the first excited state of the target nucleus, E_1 , results in simple elastic scattering, and the nucleus is left in its ground state. If the neutron energy is higher than E_1 , the compound nucleus can decay to E_1 or to the ground state of the target nucleus. The lowest neutron energy capable of leaving the target nucleus in its first excited state is the energy of that state. The metastable state generally is only a few hundred kilovolts above the ground state and will be one of the first levels to be excited. The selection rules which prevent the rapid decay of a metastable state may also prevent its ready excitation from the ground state directly. Since both selection rules and energy requirements must be satisfied in nuclear reactions, it may be necessary to excite a higher energy level which will decay by gamma-ray emission to the metastable state. Figure 1 shows the direct and indirect methods of producing state E_1 (assumed metastable) in a hypothetical nucleus. The method which actually produces the metastable state depends on the specific selection rules which operate.

The cross section for exciting a metastable state is the product of the cross section, σ_c , for exciting the compound nucleus and the probability, P_M , that it will decay to the metastable state.

Copyright © 2001 by John Wiley & Sons, Inc.

The first part of the paper is devoted to a review of the literature on the topic. The second part is devoted to a review of the literature on the topic. The third part is devoted to a review of the literature on the topic. The fourth part is devoted to a review of the literature on the topic. The fifth part is devoted to a review of the literature on the topic. The sixth part is devoted to a review of the literature on the topic. The seventh part is devoted to a review of the literature on the topic. The eighth part is devoted to a review of the literature on the topic. The ninth part is devoted to a review of the literature on the topic. The tenth part is devoted to a review of the literature on the topic.




Figure 1

SCHEMATIC DIAGRAM OF NEUTRON SCATTERING

Blatt and Weisskopf (B3) have derived an expression for the cross section for the formation of the compound nucleus by neutron bombardment as a function of neutron energy and the orbital angular momentum, l , of the incoming neutron relative to the target nucleus. The expression is:

$$\sigma_0(l) = (2l + 1)\pi \lambda^2 T_l$$

where

λ = wave length of the incident neutron,

T_l = transmission coefficient for a neutron through the nuclear surface.

Assuming there is no potential for neutrons outside the target nucleus, the transmission coefficient is:

$$T_l = 4 \left(\frac{x}{X} \right) v_l$$

where

$x = kR$; k = neutron wave number outside the nucleus,

$X = KR$; K = neutron wave number inside the nucleus,

R = nuclear radius,

v_l = factor depending on x and l .

Methods for computing K and v are given by Blatt and Weisskopf.

The probability for decay of the compound nucleus to a given excited state of the target nucleus may be determined by utilizing the

Watts and Helmholtz (11) have derived an expression for

the cross section for the formation of the compound nucleus by neutron bombardment as a function of neutron energy and the optical potential parameter, l , of the imaginary neutron potential in the target nucleus. The expression is

$$\sigma(l) = \pi(l + l_0)^2 \exp(-2\chi)$$

where

χ = wave number of the incident neutron,

l = transmission coefficient for a neutron through the

target nucleus,

Assuming there is no potential for neutrons outside the target nucleus, the transmission coefficient is

$$l = \frac{1}{2} \left(1 + \frac{k}{k_0} \right)$$

where

$x = kR$; k = neutron wave number outside the nucleus,

$x = k_0R$; k_0 = neutron wave number inside the nucleus,

R = nuclear radius,

v = neutron de Broglie wavelength in cm.

Methods for computing l and χ are given by Watts and Helmholtz.

The probability for decay of the compound nucleus to a given emitted state of the target nucleus can be determined by utilizing the

reciprocity theorem

$$\frac{\sigma(A \rightarrow B)}{(\chi_A)^2} = \frac{\sigma(B \rightarrow A)}{(\chi_B)^2}$$

which relates the cross sections for transitions in both directions between two states. Considering neutron emission as the only method of decay, the probability of the compound nuclei decaying to a state having energy E_1 above the ground state of the target nucleus compared to the n other states having energies E_i above the ground state is:

$$\gamma_1 = \frac{(E - E_1) \sigma_c(E_1, l)}{\sum_{i=1}^{i=n} (E - E_i) \sigma_c(E_i, l)}$$

where

$\sigma_c(E_i, l)$ = cross section for formation of the compound nucleus from level i ,

E = energy available for decay of the compound nucleus to the ground state of the target nucleus.

The cross section for the formation of the metastable state cannot be accurately computed unless the cross sections for the formation of the compound nucleus from all possible levels in the target nucleus are known. The calculation outlined above applies specifically to the case where the metastable state is formed only by direct decay from the compound nucleus. In many cases the compound nucleus will decay to an intermediate excited state, which then decays to the metastable state. When this condition

Probability theory

$$\frac{\partial \log L}{\partial \theta} = \frac{\sum_{i=1}^n \frac{\partial \log f(x_i; \theta)}{\partial \theta}}{n}$$

which relates the score function to both observations
between two states. The score function is the only
of state, the probability of the random variable being in a state
having mean μ shows the score of the random variable
as the random variable being in a state μ shows the score of

$$\frac{\partial \log L}{\partial \theta} = \frac{\sum_{i=1}^n \frac{\partial \log f(x_i; \theta)}{\partial \theta}}{n}$$

$$\frac{\partial \log L}{\partial \theta} = \frac{\sum_{i=1}^n \frac{\partial \log f(x_i; \theta)}{\partial \theta}}{n}$$

θ = energy available for each of the random variables in the
ground state of the input system.
The score function for the transition of the random variable is
randomly sampled when the score function for the transition of the
random variable from all possible levels is the input system are known.
The calculation of the score function for the transition of the
random variable is known only if the input system is known.
In many cases the random variable will be known to an intermediate value
state, which then depends on the random variable. This is the condition

exists, appropriate corrections, involving probabilities of gamma-ray emission, must be applied to the above calculation.

Neutron Versus Photon Excitation of Metastable States

During the process of inelastic scattering, a target nucleus may absorb from any incident neutron the energy required to raise itself to an excited state, providing the neutron possesses kinetic energy equal to or greater than the energy of the particular excited state. However, for the case of photons, production of an excited state in a target nucleus involves line absorption of only photons of the same energy as that of the excited state.

When a metastable state is produced in a nucleus, the exciting medium must carry in the angular momentum required to change the spin of the nucleus from that of the ground state to that of the metastable state. Direct production of a metastable state by quantum excitation is very improbable because the same selection rules which insure the long lifetime of a metastable state also prevent the nucleus from absorbing radiation corresponding to a transition from the ground state to the metastable state. Under such a limitation, metastable states, produced by practical intensities of photon bombardment, must be formed in two steps. A highly excited state of the nucleus is produced, which then decays by gamma-ray transitions. A fraction of these decays is to the metastable state.

Neutrons have a higher probability than quanta of carrying into the target nucleus the 3 or 4 units of angular momentum usually

required to excite the metastable state directly from the ground state. Ebel (31), using monoenergetic fast neutrons as bombarding particles, was able to excite the metastable state directly from the ground state in Au¹⁹⁷, but not in In¹¹⁵. Of course, neutrons have a greater probability of carrying only one or two units of angular momentum into a target nucleus and, therefore, may also excite those higher energy levels which are excited directly from the ground state by photon bombardment.

[illegible]

III. EXPERIMENTAL EQUIPMENT

Source of Neutrons

The $\text{Li}^7(\text{p}, \text{n})\text{Be}^7$ reaction, which gives a relatively high yield of nearly monoenergetic neutrons, was used as a source of fast neutrons for this experiment. If neutrons of known and constant energy are to be obtained from this reaction, the energy of the protons must be accurately known and must remain constant throughout individual excitation runs. Protons with the required characteristics were obtained from the W.I.T. Rockefeller electrostatic generator.

The $\text{Li}^7(\text{p}, \text{n})$ reaction is endoergic, having a Q value of -1.63 Mev. At its threshold, 1.88 Mev, it supplies 0.029 -Mev neutrons in the forward direction.

The target was prepared by evaporating metallic lithium onto a tantalum backing. To insure a uniform target thickness and to prevent boiling off of the lithium during generator operation, the tantalum backing was located eccentric to both the lithium furnace and the proton beam and was continuously rotated during target preparation or proton bombardment. The vacuum, which is maintained in the accelerator tube and deflection chamber, prevents the lithium from being oxidized. Target thicknesses were determined by Taschek and Hemmendinger's (T2) method of assuming the thickness in Kev to equal the energy increment between the neutron threshold and the

ALL INFORMATION CONTAINED

HEREIN IS UNCLASSIFIED

DATE 11/19/2013 BY 60322

DATE 11/19/2013 BY 60322

DATE 11/19/2013 BY 60322

DATE 11/19/2013 BY 60322

DATE 11/19/2013 BY 60322

DATE 11/19/2013 BY 60322

DATE 11/19/2013 BY 60322

DATE 11/19/2013 BY 60322

DATE 11/19/2013 BY 60322

DATE 11/19/2013 BY 60322

DATE 11/19/2013 BY 60322

DATE 11/19/2013 BY 60322

DATE 11/19/2013 BY 60322

DATE 11/19/2013 BY 60322

DATE 11/19/2013 BY 60322

DATE 11/19/2013 BY 60322

DATE 11/19/2013 BY 60322

DATE 11/19/2013 BY 60322

DATE 11/19/2013 BY 60322

DATE 11/19/2013 BY 60322

first (geometrical) peak of the yield curve for zero degrees. The tantalum backing stops all the protons, but does not react with neutrons.

Rockefeller Generator

The unanalyzed beam of charged particles consists of protons and diatomic and triatomic hydrogen ions. The accelerating tube is kept under vacuum, so that the ions suffer very few collisions and are nearly monoenergetic. The voltage across the terminals of the generator determines the energy of the beam. After passing down the accelerating tube, the proton beam is bent through 90 degrees by an analyzing magnetic field. The beam is focused at the top of the tube by a lens of adjustable voltage. Its cross section is defined at the bottom of the tube by two pairs of crossed adjustable slits. The details of the principle, operation, and use of the machine have been well covered in the literature (J1, W6, P1).

A nuclear-resonance method is used for controlling the field of the analyzing magnet. This allows fine control and accurate measurement of the energy of the proton beam. The frequency of the nuclear resonance is measured and is related to the energy of the protons in the beam by the expression (Schoenfeld and Duborg (34), Radden (K3)):

$$E = hf^2,$$

where

E = proton energy,

f = frequency of nuclear resonance.

Livingston (13) gives the total uncertainty in the proton energy as ± 5 Kev.

The neutron energy may be calculated using the measured value of proton energy and the laws of conservation of mass-energy and momentum. A table of neutron energies corresponding to various proton energies has been calculated in this manner by Willard (W6) and was used by the author for this investigation.

Measurement of Neutron Field

The field of neutrons was measured with a long counter similar in design to that of Hanson and McKibben (H4). This counter is shown in Figure 2. The cadmium shield prevents neutrons of very low energy (multiply scattered from the floor, walls, etc.) from entering the paraffin, and, therefore, most of the neutrons which reach the central BF_3 counter have only thermal energy. Experiments have proved that these characteristics of construction, together with the correct combination of physical dimensions, make a counter of this type equally sensitive to all neutrons with energies between 0.5 and 2.5 Mev (H4, W6). The counter used during this investigation decreases in relative sensitivity for neutrons with energies below 0.5 Mev, as shown in Figure 3 (W6).

The associated electronic equipment consisted of:

- (a) Regulated high-voltage supply (E. I. T. Servomechanisms Laboratory, Electronic Nuclear Instrumentation Project).
- (b) Model 100 preamplifier.




Figure 2
LONG COUNTER




Figure 3
RESPONSE OF THE LONG COUNTER TO MONOENERGETIC NEUTRONS

(c) Model 100 amplifier.

(d) Model 105 Atomic Instrument Company decade scaler.

Items (b) and (c) are described in detail by Elmore and Sands (12).

Scintillation Counter

The decay of the metastable state was detected by means of a scintillation counter. The counter consisted of an RCA type 5819 photomultiplier tube with an anthracene crystal cemented to its window with Canada balsam. The crystal was approximately 1 1/2 inches in diameter and 5/16 inch thick. In order to increase the light reflection inside the crystal, its exposed sides were covered with 0.00025-inch aluminum foil. A 0.04-inch mu metal shield, for the purpose of reducing stray magnetic fields, surrounded the sides of the tube and the crystal. The entire counter was encased in a 3-inch lead shield, which reduced the background due to stray radiation. The lead shield was also light-tight. A brass slider, extending into the lead shield, permitted rapid and convenient changing of experimental samples. The slider contained a circular cavity, for receiving 3-cm. diameter foils or disks, which positioned the sample in line with the axis of the tube and about 3 mm. from the crystal.

The output of the phototube is extremely sensitive, even to very small fluctuations in input voltage. Therefore, if comparable results are to be reproduced over a period of time, voltage stability is necessary throughout the electronic setup associated with the scintillation counter. Sola constant-voltage transformers were used as

449th From 501 to 510 (n)

...and the

[illegible]

1999, 2000, 2001, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022, 2023, 2024, 2025, 2026, 2027, 2028, 2029, 2030, 2031, 2032, 2033, 2034, 2035, 2036, 2037, 2038, 2039, 2040, 2041, 2042, 2043, 2044, 2045, 2046, 2047, 2048, 2049, 2050, 2051, 2052, 2053, 2054, 2055, 2056, 2057, 2058, 2059, 2060, 2061, 2062, 2063, 2064, 2065, 2066, 2067, 2068, 2069, 2070, 2071, 2072, 2073, 2074, 2075, 2076, 2077, 2078, 2079, 2080, 2081, 2082, 2083, 2084, 2085, 2086, 2087, 2088, 2089, 2090, 2091, 2092, 2093, 2094, 2095, 2096, 2097, 2098, 2099, 2100, 2101, 2102, 2103, 2104, 2105, 2106, 2107, 2108, 2109, 2110, 2111, 2112, 2113, 2114, 2115, 2116, 2117, 2118, 2119, 2120, 2121, 2122, 2123, 2124, 2125, 2126, 2127, 2128, 2129, 2130, 2131, 2132, 2133, 2134, 2135, 2136, 2137, 2138, 2139, 2140, 2141, 2142, 2143, 2144, 2145, 2146, 2147, 2148, 2149, 2150, 2151, 2152, 2153, 2154, 2155, 2156, 2157, 2158, 2159, 2160, 2161, 2162, 2163, 2164, 2165, 2166, 2167, 2168, 2169, 2170, 2171, 2172, 2173, 2174, 2175, 2176, 2177, 2178, 2179, 2180, 2181, 2182, 2183, 2184, 2185, 2186, 2187, 2188, 2189, 2190, 2191, 2192, 2193, 2194, 2195, 2196, 2197, 2198, 2199, 2200, 2201, 2202, 2203, 2204, 2205, 2206, 2207, 2208, 2209, 2210, 2211, 2212, 2213, 2214, 2215, 2216, 2217, 2218, 2219, 2220, 2221, 2222, 2223, 2224, 2225, 2226, 2227, 2228, 2229, 2230, 2231, 2232, 2233, 2234, 2235, 2236, 2237, 2238, 2239, 2240, 2241, 2242, 2243, 2244, 2245, 2246, 2247, 2248, 2249, 2250, 2251, 2252, 2253, 2254, 2255, 2256, 2257, 2258, 2259, 2260, 2261, 2262, 2263, 2264, 2265, 2266, 2267, 2268, 2269, 2270, 2271, 2272, 2273, 2274, 2275, 2276, 2277, 2278, 2279, 2280, 2281, 2282, 2283, 2284, 2285, 2286, 2287, 2288, 2289, 2290, 2291, 2292, 2293, 2294, 2295, 2296, 2297, 2298, 2299, 2300, 2301, 2302, 2303, 2304, 2305, 2306, 2307, 2308, 2309, 2310, 2311, 2312, 2313, 2314, 2315, 2316, 2317, 2318, 2319, 2320, 2321, 2322, 2323, 2324, 2325, 2326, 2327, 2328, 2329, 2330, 2331, 2332, 2333, 2334, 2335, 2336, 2337, 2338, 2339, 2340, 2341, 2342, 2343, 2344, 2345, 2346, 2347, 2348, 2349, 2350, 2351, 2352, 2353, 2354, 2355, 2356, 2357, 2358, 2359, 2360, 2361, 2362, 2363, 2364, 2365, 2366, 2367, 2368, 2369, 2370, 2371, 2372, 2373, 2374, 2375, 2376, 2377, 2378, 2379, 2380, 2381, 2382, 2383, 2384, 2385, 2386, 2387, 2388, 2389, 2390, 2391, 2392, 2393, 2394, 2395, 2396, 2397, 2398, 2399, 2400, 2401, 2402, 2403, 2404, 2405, 2406, 2407, 2408, 2409, 2410, 2411, 2412, 2413, 2414, 2415, 2416, 2417, 2418, 2419, 2420, 2421, 2422, 2423, 2424, 2425, 2426, 2427, 2428, 2429, 2430, 2431, 2432, 2433, 2434, 2435, 2436, 2437, 2438, 2439, 2440, 2441, 2442, 2443, 2444, 2445, 2446, 2447, 2448, 2449, 2450, 2451, 2452, 2453, 2454, 2455, 2456, 2457, 2458, 2459, 2460, 2461, 2462, 2463, 2464, 2465, 2466, 2467, 2468, 2469, 2470, 2471, 2472, 2473, 2474, 2475, 2476, 2477, 2478, 2479, 2480, 2481, 2482, 2483, 2484, 2485, 2486, 2487, 2488, 2489, 2490, 2491, 2492, 2493, 2494, 2495, 2496, 2497, 2498, 2499, 2500, 2501, 2502, 2503, 2504, 2505, 2506, 2507, 2508, 2509, 2510, 2511, 2512, 2513, 2514, 2515, 2516, 2517, 2518, 2519, 2520, 2521, 2522, 2523, 2524, 2525, 2526, 2527, 2528, 2529, 2530, 2531, 2532, 2533, 2534, 2535, 2536, 2537, 2538, 2539, 2540, 2541, 2542, 2543, 2544, 2545, 2546, 2547, 2548, 2549, 2550, 2551, 2552, 2553, 2554, 2555, 2556, 2557, 2558, 2559, 2560, 2561, 2562, 2563, 2564, 2565, 2566, 2567, 2568, 2569, 2570, 2571, 2572, 2573, 2574, 2575, 2576, 2577, 2578, 2579, 2580, 2581, 2582, 2583, 2584, 2585, 2586, 2587, 2588, 2589, 2590, 2591, 2592, 2593, 2594, 2595, 2596, 2597, 2598, 2599, 2600, 2601, 2602, 2603, 2604, 2605, 2606, 2607, 2608, 2609, 2610, 2611, 2612, 2613, 2614, 2615, 2616, 2617, 2618, 2619, 2620, 2621, 2622, 2623, 2624, 2625, 2626, 2627, 2628, 2629, 2630, 2631, 2632, 2633, 2634, 2635, 2636, 2637, 2638, 2639, 2640, 2641, 2642, 2643, 2644, 2645, 2646, 2647, 2648, 2649, 2650, 2651, 2652, 2653, 2654, 2655, 2656, 2657, 2658, 2659, 2660, 2661, 2662, 2663, 2664, 2665, 2666, 2667, 2668, 2669, 2670, 2671, 2672, 2673, 2674, 2675, 2676, 2677, 2678, 2679, 2680, 26

[illegible]

sources of power for all electronic units, and a special voltage stabilizer was inserted between the high-voltage supply and the phototube input.

The power for the phototube was supplied by a regulated high-voltage supply of the type described in the previous section. The phototube signal was fed through a cathode-follower preamplifier into a Model 100 amplifier (E2). The output from the amplifier was fed into a Model 210 (scale-of-32) single-channel differential discriminator which had been altered in the manner described in Appendix A.

The reproducibility of results given by this equipment was established by counting the decay of radioactive standards at frequent intervals during the investigation.

Platinum-Screen Counter

A platinum-screen gamma-ray counter, which is currently used by the Radioactivity Center of the M.I.T. Physics Department for determining absolute disintegration rates, was employed to obtain the absolute rate of emission of gamma rays by a sample of Hg^{203} . The counter is similar to the platinum-screen counter investigated by Peacock (P2) during his determination of gamma-ray counter efficiencies. The standardized Hg^{203} sample was needed for estimating the efficiency of the scintillation counter.

100-443888-100

The results of the investigation are summarized in the following table:

1996, 2000, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022, 2023, 2024, 2025, 2026, 2027, 2028, 2029, 2030, 2031, 2032, 2033, 2034, 2035, 2036, 2037, 2038, 2039, 2040, 2041, 2042, 2043, 2044, 2045, 2046, 2047, 2048, 2049, 2050, 2051, 2052, 2053, 2054, 2055, 2056, 2057, 2058, 2059, 2060, 2061, 2062, 2063, 2064, 2065, 2066, 2067, 2068, 2069, 2070, 2071, 2072, 2073, 2074, 2075, 2076, 2077, 2078, 2079, 2080, 2081, 2082, 2083, 2084, 2085, 2086, 2087, 2088, 2089, 2090, 2091, 2092, 2093, 2094, 2095, 2096, 2097, 2098, 2099, 2100, 2101, 2102, 2103, 2104, 2105, 2106, 2107, 2108, 2109, 2110, 2111, 2112, 2113, 2114, 2115, 2116, 2117, 2118, 2119, 2120, 2121, 2122, 2123, 2124, 2125, 2126, 2127, 2128, 2129, 2130, 2131, 2132, 2133, 2134, 2135, 2136, 2137, 2138, 2139, 2140, 2141, 2142, 2143, 2144, 2145, 2146, 2147, 2148, 2149, 2150, 2151, 2152, 2153, 2154, 2155, 2156, 2157, 2158, 2159, 2160, 2161, 2162, 2163, 2164, 2165, 2166, 2167, 2168, 2169, 2170, 2171, 2172, 2173, 2174, 2175, 2176, 2177, 2178, 2179, 2180, 2181, 2182, 2183, 2184, 2185, 2186, 2187, 2188, 2189, 2190, 2191, 2192, 2193, 2194, 2195, 2196, 2197, 2198, 2199, 2200, 2201, 2202, 2203, 2204, 2205, 2206, 2207, 2208, 2209, 2210, 2211, 2212, 2213, 2214, 2215, 2216, 2217, 2218, 2219, 2220, 2221, 2222, 2223, 2224, 2225, 2226, 2227, 2228, 2229, 2230, 2231, 2232, 2233, 2234, 2235, 2236, 2237, 2238, 2239, 2240, 2241, 2242, 2243, 2244, 2245, 2246, 2247, 2248, 2249, 2250, 2251, 2252, 2253, 2254, 2255, 2256, 2257, 2258, 2259, 2260, 2261, 2262, 2263, 2264, 2265, 2266, 2267, 2268, 2269, 2270, 2271, 2272, 2273, 2274, 2275, 2276, 2277, 2278, 2279, 2280, 2281, 2282, 2283, 2284, 2285, 2286, 2287, 2288, 2289, 2290, 2291, 2292, 2293, 2294, 2295, 2296, 2297, 2298, 2299, 2300, 2301, 2302, 2303, 2304, 2305, 2306, 2307, 2308, 2309, 2310, 2311, 2312, 2313, 2314, 2315, 2316, 2317, 2318, 2319, 2320, 2321, 2322, 2323, 2324, 2325, 2326, 2327, 2328, 2329, 2330, 2331, 2332, 2333, 2334, 2335, 2336, 2337, 2338, 2339, 2340, 2341, 2342, 2343, 2344, 2345, 2346, 2347, 2348, 2349, 2350, 2351, 2352, 2353, 2354, 2355, 2356, 2357, 2358, 2359, 2360, 2361, 2362, 2363, 2364, 2365, 2366, 2367, 2368, 2369, 2370, 2371, 2372, 2373, 2374, 2375, 2376, 2377, 2378, 2379, 2380, 2381, 2382, 2383, 2384, 2385, 2386, 2387, 2388, 2389, 2390, 2391, 2392, 2393, 2394, 2395, 2396, 2397, 2398, 2399, 2400, 2401, 2402, 2403, 2404, 2405, 2406, 2407, 2408, 2409, 2410, 2411, 2412, 2413, 2414, 2415, 2416, 2417, 2418, 2419, 2420, 2421, 2422, 2423, 2424, 2425, 2426, 2427, 2428, 2429, 2430, 2431, 2432, 2433, 2434, 2435, 2436, 2437, 2438, 2439, 2440, 2441, 2442, 2443, 2444, 2445, 2446, 2447, 2448, 2449, 2450, 2451, 2452, 2453, 2454, 2455, 2456, 2457, 2458, 2459, 2460, 2461, 2462, 2463, 2464, 2465, 2466, 2467, 2468, 2469, 2470, 2471, 2472, 2473, 2474, 2475, 2476, 2477, 2478, 2479, 2480, 2481, 2482, 2483, 2484, 2485, 2486, 2487, 2488, 2489, 2490, 2491, 2492, 2493, 2494, 2495, 2496, 2497, 2498, 2499, 2500, 2501, 2502, 2503, 2504, 2505, 2506, 2507, 2508, 2509, 2510, 2511, 2512, 2513, 2514, 2515, 2516, 2517, 2518, 2519, 2520, 2521, 2522, 2523, 2524, 2525, 2526, 2527, 2528, 2529, 2530, 2531, 2532, 2533, 2534, 2535, 2536, 2537, 2538, 2539, 2540, 2541, 2542, 2543, 2544, 2545, 2546, 2547, 2548, 2549, 2550, 2551, 2552, 2553, 2554, 2555, 2556, 2557, 2558, 2559, 2560, 2561, 2562, 2563, 2564, 2565, 2566, 2567, 2568, 2569, 2570, 2571, 2572, 2573, 2574, 2575, 2576, 2577, 2578, 2579, 2580, 2581, 2582, 2583, 2584, 2585, 2586, 2587, 2588, 2589, 2590, 2591, 2592, 2593, 2594, 2595, 2596, 2597, 2598, 2599, 2600, 2601, 2602, 2603, 2604, 2605, 2606, 2607, 2608, 2609, 2610, 2611, 2612, 2613, 2614, 2615, 2616, 2617, 2618, 2619, 2620, 2621, 2622, 2623, 2624, 2625, 2626, 2627, 2628, 2629, 2630, 2631, 2632, 2633, 2634, 2635, 2636, 2637, 2638, 2639, 2640, 2641, 2642, 2643, 2644, 2645, 2646, 2647, 2648, 2649, 2650, 2651, 2652, 2653, 2654, 2655, 2656, 2657, 2658, 2659, 2660, 2661, 2662, 2663, 2664, 2665, 2666, 2667, 2668, 2669, 2670, 2671, 2672, 2673, 2674, 2675, 2676, 2677, 2678, 2679, 2680, 2681, 26

[illegible]

IV. INVESTIGATION OF Cd^{111} Historical Background of the Excitation of Metastable Cd^{111}

Mattuch, the Austrian physicist, has postulated two rules regarding the existence of metastable states:

- (a) Isomerism does not occur in nuclei having an even number of protons and an even number of neutrons.
- (b) Isomerism occurs only in nuclei which have ground-state spins of $1/2$ or $3/2$ or greater.

Pb^{204} , an even-even nucleus, is usually reported as having a 68-minute metastable state (not definitely established as isomeric) and appears to violate rule (a). Au^{197} , having a ground-state spin of $3/2$ and a 7.5-second metastable state, violates rule (b). However, such rules should not be expected to be without exception, and they at least suggest possible isomeric nuclei for a given mass number.

Cd^{111} , with a ground-state spin of $1/2$, conforms with both of Mattuch's rules. Therefore, it was not surprising that Wiedenbeck (W7), in 1944, reported the existence of a 45.6-minute metastable state in Cd^{111} . Wiedenbeck erroneously determined this energy to be 0.195 Mev above the ground state. Hole (H1), in 1947, determined spectrographically that the energy of the metastable state was 0.361 Mev above the ground state and that it decays by gamma emission to a level 0.235 Mev above the ground state, which then decayed to the ground state by gamma emission. The currently accepted values for these levels, 0.396 Mev and 0.247 Mev, were determined by Helmholtz, Hayward, and McGinnis (H5) with

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

(a) THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

(b) THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

a magnetic lens spectrograph. Goldhaber and Nuelhouse (G2), by observing the (n,γ) reaction in Cd^{110} , have confirmed that the 48.6-minute metastable state exists in Cd^{111} . Deutsch (D2, D3) subsequently determined that the 0.247-Mev excited level decayed with a half-life of 8×10^{-8} second, and this level has been reclassified as a second metastable state in Cd^{111} . Therefore, this isotope is unique because it is the only stable nucleus currently known to have more than one metastable state.

Energy Levels in Cd^{111}

Figure 4 is an energy level diagram for Cd^{111} . It contains all of the currently published information (W1) about the decay of the excited states of the nucleus and about the decay of neighboring isotopes to Cd^{111} by beta emission and K-capture. The presence of the four highest excited levels was indicated by Wiedenbeck's (W7) excitation curves for the 48.6-minute metastable state. He obtained curves for both X-ray and electron bombardment of Cd^{111} . The threshold energy observed by Wiedenbeck for production of the metastable state was 1.25 Mev for both types of excitation.

The theoretical half-lives against decay by gamma-ray emission of several different multipole orders for both metastable states of Cd^{111} have been calculated by the author using various formulas based on the Wigner-Eisenberg hypothesis. The half-lives obtained by these calculations, both uncorrected and corrected for internal conversion, are presented in Table I.

a solution was prepared, which was then used for the

analysis of the (12) series at 25°C. (see Table I).

The results of the analysis are given in Table I. The results

show that the 0.1M solution of the polymer in a solution of

0.1M NaCl, and the 0.1M solution of the polymer in a solution of

0.1M NaCl, are identical. This shows that the polymer is

in the form of a single molecule in the form of a polymer.

Table I.

Table I.

Figure 1 is a graph of the log of the ratio of the

log of the ratio of the log of the ratio of the log of the

log of the ratio of the log of the ratio of the log of the

log of the ratio of the log of the ratio of the log of the

log of the ratio of the log of the ratio of the log of the

log of the ratio of the log of the ratio of the log of the

log of the ratio of the log of the ratio of the log of the

log of the ratio of the log of the ratio of the log of the

log of the ratio of the log of the ratio of the log of the

log of the ratio of the log of the ratio of the log of the

log of the ratio of the log of the ratio of the log of the

log of the ratio of the log of the ratio of the log of the

log of the ratio of the log of the ratio of the log of the

log of the ratio of the log of the ratio of the log of the

log of the ratio of the log of the ratio of the log of the




Figure 4

ENERGY LEVEL DIAGRAM FOR Cd^{113}

TABLE I

Cd^{111} $E_\gamma = 0.149 \text{ Mev}$ $\lambda_a = 12$ $\tau_{1/2 \text{ exp.}} = 48.6 \text{ min.}$ Half-life units = minutes

L	Theoretical			Corrected for Internal Conversion			
	Bethe (S)	Lowen (M)	Berthelot (M)	Neon (M)	Bethe	Lowen	Berthelot
3	4.82×10^{-3}	9.20×10^{-6}	1.15×10^{-6}	2.99×10^{-5}	3.71×10^{-4}	7.07×10^{-7}	8.86×10^{-8}
4	1.39×10^{-4}	1.79×10^{-1}	3.42×10^{-1}	5.75×10^{-1}	1.07×10^{-3}	1.36	2.63×10^{-2}
5	6.98×10^{-10}	6.07×10^{-7}	1.92×10^{-5}	very long	5.37×10^{-9}	4.67×10^{-6}	1.48×10^{-4}
							very long

Cd^{111} $E_\gamma = 0.247 \text{ Mev}$ $\lambda_a = 0.1$ $\tau_{1/2 \text{ exp.}} = 8 \times 10^{-8} \text{ sec.}$ Half-life units = seconds

2	2.48×10^{-8}	5.97×10^{-11}	---	1.16×10^{-10}	2.26×10^{-8}	5.42×10^{-11}	---
3	2.20×10^{-2}	2.65×10^{-5}	3.00×10^{-6}	6.95×10^{-5}	1.00×10^{-2}	2.41×10^{-5}	2.72×10^{-6}
							1.05×10^{-10}
							6.32×10^{-5}

λ_a = Internal conversion coefficient.

**Order of electric multipole = Λ , order of magnetic multipole = $\Lambda - 1$ (Magnetic $2^{\Lambda} - \text{pole}$ radiation has approximately the same probability of occurring as electric $2^{\Lambda} + 1 - \text{pole}$ radiation.)

A comparison of the experimentally determined 43.6-minute half-life of the 0.396-Mev level with the theoretical half-lives corrected for internal conversion indicates quite clearly that the transition from this level to the 0.247-Mev level is by electric 2^h -pole or magnetic octupole radiation ($\Lambda = 4$). The assignment of $\Lambda = 4$ to the 0.149-Mev transition coincides with the multipole order designated by Helmholtz, Hayward, and McGinnis (H5) and by Axel and Dancoff (A1) for this transition. A similar comparison for the transition from the 0.247-Mev level to the ground state ($T_{1/2 \text{ exp.}} = 5 \times 10^{-8}$ second) indicates that Λ is 2 or 3, but does not allow a reliable assignment of a specific value to .

A knowledge of the half-life alone does not allow determination of whether a gamma-ray transition involves an electric or a magnetic multipole, but a comparison of experimentally obtained K or L conversion coefficients or the K/L ratio of the coefficients with theoretically calculated values of the same quantities may facilitate such a determination. Table II gives currently accepted experimental and calculated theoretical data concerning internal conversion for probable values of Λ for transitions from both metastable states in Cd^{111} . The theoretical values were calculated from formulas developed by Dancoff and Morrison (D1) or were obtained from tables or graphs, based on these formulas, prepared by Hebb and Nelson (H6). Table II indicates with reliable certainty that the transition from the 43.6-minute level is by electric 2^h -pole radiation (the same assignment has been made by Helmholtz et al. (H5)). If the transition from the 5×10^{-8} -second level is of

TABLE II

DATA ON INTERNAL CONVERSION OF Cd^{111}

Experimental			Theoretical	
			Electric	Magnetic
$E_\gamma = 0.149 \text{ Mev}$ $\Delta = 4.00$	α_K	7.72	5.63	3.40
	α_L	4.28	6.05	0.66
	α_K/α_L	1.80	0.906	5.08
$E_\gamma = 0.247 \text{ Mev}$ $\Delta = 2.00$	α_K	0.09	0.0394	0.015
	α_L	0.01	0.00584	0.0021
	α_K/α_L	10.00	6.74	7.22
$E_\gamma = 0.247 \text{ Mev}$ $\Delta = 3.00$	α_K	0.09	0.117	0.077
	α_L	0.01	0.0165	0.0112
	α_K/α_L	10.00	6.31	6.90

 α_K = K shell internal conversion coefficient. α_L = L shell internal conversion coefficient.

TABLE II

TABLE II. LITERATURE DATA ON THE
 RATE OF DECOMPOSITION OF H_2O_2

Temperature, $^{\circ}\text{C}$ Rate constant, min^{-1}

Temperature, $^{\circ}\text{C}$	Rate constant, min^{-1}	Rate constant, min^{-1}	Rate constant, min^{-1}	Rate constant, min^{-1}
20.0	0.0001	0.0001	0.0001	0.0001
25.0	0.0002	0.0002	0.0002	0.0002
30.0	0.0004	0.0004	0.0004	0.0004
35.0	0.0008	0.0008	0.0008	0.0008
40.0	0.0016	0.0016	0.0016	0.0016
45.0	0.0032	0.0032	0.0032	0.0032
50.0	0.0064	0.0064	0.0064	0.0064
55.0	0.0128	0.0128	0.0128	0.0128
60.0	0.0256	0.0256	0.0256	0.0256
65.0	0.0512	0.0512	0.0512	0.0512
70.0	0.1024	0.1024	0.1024	0.1024
75.0	0.2048	0.2048	0.2048	0.2048
80.0	0.4096	0.4096	0.4096	0.4096
85.0	0.8192	0.8192	0.8192	0.8192
90.0	1.6384	1.6384	1.6384	1.6384
95.0	3.2768	3.2768	3.2768	3.2768
100.0	6.5536	6.5536	6.5536	6.5536

TABLE II. LITERATURE DATA ON THE
 RATE OF DECOMPOSITION OF H_2O_2

TABLE II. LITERATURE DATA ON THE
 RATE OF DECOMPOSITION OF H_2O_2

the $\Delta = 3$ type, Table II indicates that it is probably by magnetic quadrupole radiation, but if $\Delta = 2$, the table fails to indicate specifically whether the transition is by electric quadrupole or magnetic dipole radiation. The difficulty of assigning the correct multipole order to this transition is emphasized by the contradictory reports in the recent literature concerning this transition (D2, E1, E5).

A metastable state is defined as any excited state having a measurable lifetime. Weizsacker showed that assigning an angular momentum to the metastable state differing by 3 to 5 units from the angular momentum of the state of lower energy to which the metastable state decays by gamma-ray emission justifies theoretically its observable lifetime. Instrumentation and measuring techniques have improved so much since Weizsacker suggested his hypothesis that decays of excited states with very short lifetimes are now detectable. Therefore, isomers which have recently been (6×10^{-6} second Cd^{111}) or in the near future will be reclassified as metastable may have angular momentum differences less than 3, because of their very short lifetimes.

Figure 5 shows all the possible combinations of angular momentums for the ground and metastable states of Cd^{111} for those decay schemes which are indicated by a comparison of theory and experiment.

$$\Lambda_{2-1}=4 \quad \Lambda_{1-0}=2 \quad E_{2-1}=0.149\text{Mev} \quad 2^4\text{-pole electric}$$

$$E_{1-0}=0.247\text{Mev} \quad 2^2\text{-pole electric}$$

	Energy	Parity	Possible Values of Angular Momentum											
I_2	0.396	+	$\frac{*13}{2}$	$\frac{11}{2}$	$\frac{9}{2}$	$\frac{7}{2}$	$\frac{5}{2}$	$\frac{3}{2}$	$\frac{11}{2}$	$\frac{9}{2}$	$\frac{7}{2}$	$\frac{5}{2}$		
I_1	0.247	+	$\frac{5}{2}$	$\frac{5}{2}$	$\frac{5}{2}$	$\frac{5}{2}$	$\frac{5}{2}$	$\frac{5}{2}$	$\frac{3}{2}$	$\frac{3}{2}$	$\frac{3}{2}$	$\frac{3}{2}$		
I_0	0	+	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$		

$$\Lambda_{2-1}=4 \quad \Lambda_{1-0}=2 \quad E_{2-1}=0.149\text{Mev} \quad 2^4\text{-pole electric}$$

$$E_{1-0}=0.247\text{Mev} \quad 2^1\text{-pole magnetic}$$

	Energy	Parity	Possible Values of Angular Momentum											
I_2	0.396	+	$\frac{*11}{2}$	$\frac{9}{2}$	$\frac{7}{2}$	$\frac{5}{2}$			$\frac{9}{2}$	$\frac{7}{2}$				
I_1	0.247	+	$\frac{3}{2}$	$\frac{3}{2}$	$\frac{3}{2}$	$\frac{3}{2}$			$\frac{1}{2}$	$\frac{1}{2}$				
I_0	0	+	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$			$\frac{1}{2}$	$\frac{1}{2}$				

$$\Lambda_{2-1}=4 \quad \Lambda_{1-0}=3 \quad E_{2-1}=0.149\text{Mev} \quad 2^4\text{-pole electric}$$

$$E_{1-0}=0.247\text{Mev} \quad 2^2\text{-pole magnetic}$$

	Energy	Parity	Possible Values of Angular Momentum											
I_2	0.396	+	$\frac{*13}{2}$	$\frac{11}{2}$	$\frac{9}{2}$	$\frac{7}{2}$	$\frac{5}{2}$	$\frac{3}{2}$	$\frac{11}{2}$	$\frac{9}{2}$	$\frac{7}{2}$	$\frac{5}{2}$		
I_1	0.247	+	$\frac{5}{2}$	$\frac{5}{2}$	$\frac{5}{2}$	$\frac{5}{2}$	$\frac{5}{2}$	$\frac{5}{2}$	$\frac{3}{2}$	$\frac{3}{2}$	$\frac{3}{2}$	$\frac{3}{2}$		
I_0	0	+	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$		

* Most Probable Combination.

Figure 5

Method of Obtaining the Uncorrected Excitation Curve

Whether a detectable amount of activity of a metastable state may be excited or not depends on the cross section for the formation of the metastable state by the particular type of excitation process employed. The cross section for the formation of a metastable state by the inelastic scattering of neutrons is usually very small. The ease with which the decay of a metastable state may be detected depends on the energies of the gamma rays and electrons emitted during the decay and on the sensitivity of the detection equipment. Therefore, high backgrounds or substantial amounts of interfering radioactivities cannot be tolerated. Possible sources of background during this investigation were stray radiation from nuclear machines, sources, etc., cosmic radiation, light leaks, and thermal emission in the photomultiplier tube itself. Prior to the actual experiment, considerable time was spent in reducing to a minimum the amount of background available to the crystal and the phototube of the scintillation counter. Interfering radioactive decays, which may accompany the decay of the metastable state, result from nuclear reactions initiated in the impurities or other isotopes present in the sample by the bombarding neutrons.

The differential discriminator was especially applicable for use as a counter in this experiment because it counts only activity of the particular energies to which its window is sensitive. It

was five times less sensitive to background than an Atomic Instrument decade scaler, when both instruments were counting the same background simultaneously.

The experimental samples were made from high-purity commercial cadmium sheet manufactured by the Belmont Smelting and Refining Company, of Newark, New Jersey. A spectroscopic analysis of this cadmium indicated that it contained about 0.1 per cent copper and 0.1 per cent lead, but no impurities were present in sufficient quantities to interfere with detecting the decay of the metastable state. Unfortunately, the isotope, Cd^{111} , is only 12.8 per cent abundant in natural cadmium. Table III contains data on all the stable isotopes of cadmium and gives the nuclear reactions which can occur when these isotopes are subjected to bombardment by neutrons of the energies used in this experiment. Decay of Cd^{107} and Cd^{109} is not important because the per cent abundance of Cd^{106} and Cd^{108} in natural cadmium is so low. No reaction between Cd^{112} and neutrons of moderate or lesser energies has been observed. Cd^{112} and Cd^{114} are stable nuclides, so no further nuclear reactions result from neutron capture by Cd^{111} and Cd^{113} . Therefore, the only possible reactions which may interfere with detection the decay of metastable $^*\text{Cd}^{111}$ formed from stable Cd^{111} are:

- (a) The formation of metastable $^*\text{Cd}^{111}$ from capture of fast neutrons by Cd^{110} .
- (b) The formation of metastable $^*\text{Cd}^{113}$ from inelastic scattering of fast neutrons by Cd^{113} .

and the first step towards the solution of the problem is to determine the nature of the problem itself. The second step is to determine the nature of the solution itself. The third step is to determine the nature of the solution itself.

The first step is to determine the nature of the problem itself. The second step is to determine the nature of the solution itself. The third step is to determine the nature of the solution itself.

The first step is to determine the nature of the problem itself. The second step is to determine the nature of the solution itself. The third step is to determine the nature of the solution itself.

The first step is to determine the nature of the problem itself. The second step is to determine the nature of the solution itself. The third step is to determine the nature of the solution itself.

The first step is to determine the nature of the problem itself. The second step is to determine the nature of the solution itself. The third step is to determine the nature of the solution itself.

The first step is to determine the nature of the problem itself. The second step is to determine the nature of the solution itself. The third step is to determine the nature of the solution itself.

The first step is to determine the nature of the problem itself. The second step is to determine the nature of the solution itself. The third step is to determine the nature of the solution itself.

The first step is to determine the nature of the problem itself. The second step is to determine the nature of the solution itself. The third step is to determine the nature of the solution itself.

TABLE III

ISOTOPES OF CADMIUM

Per Cent Abundance	Isotope	Reaction	Half-Life	Decay Radiations
1.1	Cd^{106}	$\text{Cd}^{106} + n \rightarrow \text{Cd}^{107}$	6.7 hours	K , $\gamma(0.546)$, $\beta^+(0.32)$
0.9	Cd^{108}	$\text{Cd}^{108} + n \rightarrow \text{Cd}^{109}$	1.3 years	K
12.4	Cd^{110}	$\text{Cd}^{110} + n \rightarrow \text{Cd}^{111}$	45.6 minutes	$\gamma(0.149)$ I.T. β^- , $\gamma(0.247)$ I.T. β^-
12.8	Cd^{111}	$\text{Cd}^{111} + n \rightarrow \text{Cd}^{111} + n$	45.6 minutes	$\gamma(0.149)$ I.T. β^- , $\gamma(0.247)$ I.T. β^-
24.1	Cd^{112}			
12.3	Cd^{113}	$\text{Cd}^{113} + n \rightarrow \text{Cd}^{113} + n$	2.3 minutes	I.T. β^-
		$\text{Cd}^{113} + n \rightarrow \text{Cd}^{114}$	Stable	
26.8	Cd^{114}	$\text{Cd}^{114} + n \rightarrow \text{Cd}^{115}$	2.33 days	$\beta^-(0.6, 1.13)$, $\gamma(0.52)$
			43.0 days	$\beta^-(1.67)$, $\gamma(0.5)$
7.6	Cd^{116}	$\text{Cd}^{116} + n \rightarrow \text{Cd}^{117}$	2.83 days	$\beta^-(1.5)$

Notes: (1) Decay radiations are in energy units of Mev.
 (2) I.T. designates isomeric transition.
 (3) β^- designates internal conversion.
 (4) * designates excited state.

1993

Year	Production (in tons)	Consumption (in tons)	Stock (in tons)	Balance (in tons)	Remarks
1970	1000	800	200	200	
1971	1200	900	300	300	
1972	1500	1100	400	400	
1973	1800	1300	500	500	
1974	2000	1500	500	500	
1975	2200	1700	500	500	
1976	2500	2000	500	500	
1977	2800	2300	500	500	
1978	3000	2500	500	500	
1979	3200	2700	500	500	
1980	3500	3000	500	500	

FREE

- (c) The formation of unstable Cd^{115} and Cd^{117} from capture of fast neutrons by Cd^{114} and Cd^{116} .
- (d) The formation of $^*\text{Cd}^{111}$, Cd^{115} , or Cd^{117} by capture of thermal neutrons.

Ways of proving that these interfering reactions are unimportant or of correcting for their effect will be described in later sections of this chapter.

Figure 6 shows the experimental arrangement used for irradiating the cadmium samples. The 48.6-minute half-life of the 0.396-Mev metastable state in Cd^{111} permitted locating the counting equipment remote from the site of the irradiation. Figure 7 shows the setup of the scintillation counter and its associated electronic equipment used for detecting the decay of the 48.6-minute metastable state.

The long half-lives of some of the unstable isotopes of cadmium which result from neutron capture prevented reusing a sample until approximately two weeks after a previous irradiation. The 25 samples used during the investigation were cadmium disks 3 cm. in diameter and 1.626 mm. thick. The mass of the disks varied between 10.015 and 10.045 grams. Relatively heavy disks were used rather than light, thin foils, in order to insure the production of a significant amount of activity, even though the cross section for the formation of the metastable state was very small.

During these experimental runs which were used to obtain the excitation curve, the disk was positioned by an aluminum holder with its

Figure 6. Arrangement for irradiating the cadmium disks, showing the magnet and chamber of the Rockefeller generator, the tantalum-backed lithium target and the aluminum disk holder.

The first of these is the fact that the
 the first of these is the fact that the
 the first of these is the fact that the

the first of these is the fact that the
 the first of these is the fact that the
 the first of these is the fact that the

the first of these is the fact that the
 the first of these is the fact that the
 the first of these is the fact that the

the first of these is the fact that the
 the first of these is the fact that the
 the first of these is the fact that the

the first of these is the fact that the
 the first of these is the fact that the
 the first of these is the fact that the

the first of these is the fact that the
 the first of these is the fact that the
 the first of these is the fact that the


the first of these is the fact that the
 the first of these is the fact that the
 the first of these is the fact that the

the first of these is the fact that the
 the first of these is the fact that the
 the first of these is the fact that the



Figure 7. Setup of equipment used for detecting the decay of the metastable state, showing the high-voltage power supply, voltage stabilizer, lead-sheathed scintillation counter, cathode-follower pre-amplifier, amplifier, differential discriminator, and decade scaler.

Figure 7. Setup of equipment used for determining the decay of the metastable state, showing the high-voltage power supply, voltage divider, fast-scan oscilloscope, and oscilloscope. The pre-amplifier, amplifier, and oscilloscope are also shown.



was operated at 1050 volts, and the coarse and fine gain settings of the amplifier were 9 and 7, respectively. The operation of the amplifier in conjunction with the differential discriminator was improved by adding a one-microsecond delay line clipper to the amplifier circuit. Preliminary experiments indicated that this combination of adjustments to the counting equipment gave the maximum ratio of desired decay counts to background counts. The window of the differential discriminator was adjusted to count preferentially the decays of ^{111}Cd and to discriminate as much as possible against any other interfering decays. The actual width and location of the window were such that all pulses between 6 and 28 volts were counted. The activity produced in the sample increased as the energy of the bombarding neutrons increased, and the number of background counts varied from 50 per cent to 4 per cent of the total counts, depending on the activity of the sample. The number of scintillation-counter counts occurring during the 45-minute counting period was recorded for each sample.

The induced activity in the sample (scintillation-counter counts/long-counter counts) was plotted against the maximum energy of the neutrons traversing the sample during an irradiation. This neutron-excitation curve is shown in Figure 5. The portion of the curve below a neutron energy of 0.5 Mev has been adjusted, as indicated by Figure 3, for the decrease of long-counter sensitivity at these lower energies. The swift rise of the curve at energies just above the neutron energy associated with the lithium threshold may be due to the decrease in

was operated at 1000 volts, and the source was then kept constant at the amplifier was 5 and 1, respectively. The operation of the amplifier in conjunction with the differential amplifier was improved by using a one-dimensional relay line applied to the amplifier circuit. Preliminary experiments indicated that this combination of adjustments to the counting equipment gave the maximum ratio of desired decay counts to background counts. The window of the differential discriminator was adjusted to about 10% of the range of ^{235}U and ^{238}U to discriminate as much as possible against any other interfering decay. The signal with the location of the window was such that all pulses between 5 and 10 volts were counted. The activity produced in the sample increased as the energy of the bombarding neutron increased, and the number of background counts varied from 50 per cent to 10 per cent of the total counts, depending on the activity of the sample. The number of calibration-counter events occurring during the 15-minute counting period was recorded for each sample.

The induced activity in the sample (calibration-counter counts/total-counter counts) was plotted against the neutron energy of the neutrons traversing the sample during the irradiation. This neutron calibration curve is shown in Figure 6. The position of the curve below a neutron energy of 0.1 Mev has been adjusted, as indicated by Figure 7, for the decrease of half-counter sensitivity at those lower energies. The half-life of the curve at energies just above the neutron energy associated with the fission threshold was also in the decrease in

center in line with and perpendicular to the proton beam and at a distance of 4 cm. from the lithium target. The holder was made from aluminum because aluminum is almost completely transparent to neutrons of all energies. The disks were irradiated for periods of one hour with monoenergetic neutrons of various energies obtained from the $\text{Li}^7(p,n)$ reaction using a 60 Kev thick target on the Rockefeller generator. Fifteen minutes after the end of the irradiation, the radioactive decays occurring in the disks were counted in the scintillation counter.

If the different points on the excitation curve are to have comparative values relative to each other, the neutron flux traversing the sample must be uniform over the entire period of each irradiation. Several times during this investigation the Rockefeller generator operated unsatisfactorily and the proton beam was unstable and sporadic. Runs which were marred by lengthy interruptions of the neutron flux were of little value.

The relative number of neutrons traversing the sample for the different points on the excitation curve was determined by the long counter described in a previous section. The counter was set up one meter from the target and at zero degrees to the proton beam. The number of long-counter counts occurring during each irradiation was recorded.

Fifteen minutes after the end of the irradiation, the decays occurring in the sample were counted for a period of 45 minutes by the scintillation counter. During this investigation, photomultiplier tube

[illegible]




Figure 8
NEUTRON EXCITATION CURVE FOR ^{109}Cd UNCORRECTED

long-counter sensitivity at very low energies, a region about which Figure 3 gives no information. Alternatively, it may be due to a sharp increase in the cross section for neutron capture by some of the cadmium isotopes at very low neutron energies. Since this condition occurs at neutron energies well below the energy of the 48.6-minute metastable state, resolution of the question is not pertinent to the investigation of the metastable state. The statistics encountered in obtaining the data for the excitation curve represent an average uncertainty of 2 per cent in the value of the ordinate. The significance of the two sharp changes in the slope of the curve will be discussed after the curve has been corrected for interfering decays and for the neutron energy spread resulting from the finite target thickness, angularity, etc. The existence of these breaks in the curve was confirmed with certainty by the fact that their location was reproduced by a second set of data obtained three months after the first set.

Elimination of the Effect of Interfering Decays

The 15-minute interval between the end of the irradiation and the beginning of the 45-minute counting period allowed any 2.3-minute metastable ^{113}Cd to decay to a negligible amount before the actual counting was started and also provided the time necessary for transporting the sample from the target room of the Rockefeller generator to the location of the counting equipment.

high-sensitivity at very low energies a rather short wave-
 length is required. Alternatively, it may be that a
 sharp increase in the cross section for nuclear capture by means of
 the nucleus is observed at very low energies. Since this pro-
 bability is rather small, it is not clear that the energy of the
 nucleus is sufficient to produce the reaction. The reaction is not
 in the investigation of the reaction. The reaction is
 concerned in obtaining the data for the reaction cross section
 as a function of the energy of the incident particles.
 The significance of the two sharp changes in the slope of the curve
 will be discussed when the curve has been corrected for scattering
 effects and for the various energy losses resulting from the finite
 target thickness, geometry, etc. The values of these losses in
 the curve are indicated with arrows by the two lines. Their location
 was reproduced by a second set of data obtained from another set of
 first set.

Discussion of the Effect of Scattering Losses

The 15-minute interval between the end of the irradiation and
 the beginning of the 15-minute counting period allows any 15-minute
 counting rate to decay to a negligible amount before the actual
 counting was started and also provides the time necessary for the
 setting the sample from the target room of the detection equipment to
 the location of the counting equipment.

The shape of the fundamental excitation curve between neutron energies of 0.3 and 0.7 Mev raises a suspicion that the curve really represents the sum of several activities with different half-lives. Therefore, an irradiation identical to those described in the previous section was made, and the decay of the sample was followed until the activity was decreasing at a very slow and uniform rate. Figure 9 is a graph of the total activity of the sample against the time after the end of the irradiation plotted on semi-log coordinates. The decay graph of an activity with a single half-life should be a straight line when plotted on such coordinates. The curvature of the plot confirms the suspicion that the total activity consists of two or more separate activities with different half-lives. According to the previous section, the possible components of the total activity are 45.6-minute ^{111}Cd , 2.33-day ^{115}Cd , 43-day ^{115}Cd , and 2.8-hour ^{117}Cd .

Since no activity was observed in the sample two weeks after the irradiation, the existence of the 43-day activity was ruled out. The shape of the decay curve 20 hours after the end of the irradiation was what would be expected for the decay of the 2.33-day activity alone. This activity was subtracted graphically from the total decay curve. The slope of the right-hand portion of the resulting curve confirmed the presence of the 2.8-hour activity. Separation of the 2.8-hour activity in a similar manner revealed that the remaining activity decayed with a 45.6-minute half-life. This graphical method of half-life separation is illustrated in detail in Figure 9.

[illegible]




Figure 9 BASIC DECAY CURVE FOR IRRADIATION BY .592 Mev NEUTRONS



Sixteen additional irradiations were made throughout the range of neutron energies covered by the uncorrected excitation curve of Figure 8. Since radioactivity decays according to an exponential law, the amount of activity remaining five half-lives after its production is a negligible fraction of its initial value. Therefore, the amount of 48.6-minute activity remaining six hours after the end of the irradiation is negligible and the amount of 2.8-hour activity remaining 19 hours later is also negligible. The decay curves for each irradiation were extrapolated to zero time, and the total activity was read from them at 0, 6, and 25 hours after the end of the irradiation. Since it has been definitely established that only three separate activities exist, the knowledge of the total activity at these three times affords a means of computing mathematically the initial induced activity for each separate half-life. Next, the appropriate Bateman equation for each radioactive decay was integrated between the limits of 15 and 60 minutes. The total number of decays of each activity occurring in the 45-minute counting period was obtained by substituting the specific values of initial activity in the proper equations. Finally, this information was used to express the integral quantity of each separate activity as a fraction of the total activity observed during the 45-minute counting period. Figure 10 shows the varying shape of the total decay curves for substantially different fractions of 48.6-minute activity.




Figure 10

COMPARISON OF DECAYS FOR DIFFERENT %'s OF 48.6 MINUTE ACTIVITY

Figure 11 contains the separated excitation curves for the three different activities detected by the scintillation counter. The magnitude of each activity was obtained by multiplying the ordinates, at the proper neutron energies, from the fundamental excitation curve of Figure 8 by the fractional values calculated for each of the 17 half-life separation runs. The probable error in the ordinates of the curves of Figure 11, which is primarily due to the poor statistics at the low-activity end of the half-life separation decay curves, is estimated as approximately 5 per cent.

The possibility that some of the activity detected may be due to the capture of thermal neutrons must not be overlooked. The source of thermal neutrons could be the multiple scattering of fast neutrons from the walls and floor of the target room. Dempster (D4) has determined by mass spectrography that Cd^{113} is the isotope of cadmium mainly responsible for the absorption of thermal neutrons. Moyer, Peters, and Schmidt (M2) have also reported that their cross-section determinations for enriched cadmium isotopes indicate that absorption of thermal neutrons by all the isotopes of cadmium except Cd^{113} is negligible compared to the capture of thermal neutrons by Cd^{113} . The previous section has already pointed out that the capture of neutrons of any energy by Cd^{113} is unimportant to this investigation, because Cd^{114} is stable and no interfering decays result. The conclusion that the capture of thermal neutrons in general is unimportant was reaffirmed experimentally during this investigation. An irradiation

Figure 11 contains the computed results shown for the
three different activities defined by the relationship system. The
magnitude of each activity was obtained by multiplying the velocity
at the proper constant frequency, from the theoretical relationship curve
of Figure 2 by the theoretical values obtained for each of the 17
half-life expansion times. The probable error in the estimates of the
curves of Figure 11, which is extremely low for the first activities of
the low-activity end of the half-life expansion time curve, is
estimated as approximately 2 per cent.

The possibility that some of the activity detected may be
due to the capture of thermal neutrons must not be overlooked. The
source of thermal neutrons could be the multiple scattering of fast
neutrons from the walls and floor of the target room. Tammes (10)
has estimated by some spectroscopy that ¹¹ in the target of cal-
cium metal is responsible for the absorption of thermal neutrons. Tammes,
Tobey, and Daniels (10) have also reported that their cross-section
determination for sodium neutron induced reactions indicated
of thermal neutrons for all the isotopes of sodium except ²³Na
is negligible compared to the capture of thermal neutrons by ²³Na. The
previous section has already pointed out that the capture of neutrons
of any energy by ²³Na is important in this investigation, because
²³Na is stable and no transmuting change results. The conclusion
that the capture of thermal neutrons is known to be important was
reaffirmed experimentally during this investigation. An investigation




Figure 11
EXCITATION CURVES FOR THE SEPARATED ACTIVITIES OF
 Cd^{113} , Cd^{115} , and Cd^{117}

was made on a sandwich of three identical cadmium disks. Decay curves for an outer disk which was exposed to any background of thermal neutrons which may have existed in the target room and for the center disk which was shielded from such a background were equal within statistics.

An accurate determination of whether any of the 48.6-minute metastable Cd^{111} was being formed by the absorption of fast neutrons in Cd^{110} , instead of by the inelastic scattering of fast neutrons by Cd^{111} , was impossible. However, the relatively small amounts of 2.8-day and 2.33-hour activities observed and the dearth of the 48.6-minute activity in the vicinity of a 0.4-Mev neutron energy indicates that the cross section for this capture reaction is very small in an energy range covered by this investigation. Therefore, the effect of neutron capture by Cd^{110} has been neglected.

Correction of the 48.6-Minute Excitation Curve for Proton-Energy Spread, Target Thickness, and Angularity

The energy resolution of the abscissas of the excitation curves obtained during this investigation may be affected by three factors: fluctuations in the energy of the proton beam, the thickness of the lithium target, and the dependence of neutron energy on the angle of neutron emission relative to the direction of the proton beam.

During this experiment, the Rockefeller generator was operated with a total exit slit opening of one mm. Such an adjustment of the beam defining slits, together with other characteristics of the generator

was made on a number of days. The first was on the 1st of May. The second on the 2nd. The third on the 3rd. The fourth on the 4th. The fifth on the 5th. The sixth on the 6th. The seventh on the 7th. The eighth on the 8th. The ninth on the 9th. The tenth on the 10th. The eleventh on the 11th. The twelfth on the 12th. The thirteenth on the 13th. The fourteenth on the 14th. The fifteenth on the 15th. The sixteenth on the 16th. The seventeenth on the 17th. The eighteenth on the 18th. The nineteenth on the 19th. The twentieth on the 20th. The twenty-first on the 21st. The twenty-second on the 22nd. The twenty-third on the 23rd. The twenty-fourth on the 24th. The twenty-fifth on the 25th. The twenty-sixth on the 26th. The twenty-seventh on the 27th. The twenty-eighth on the 28th. The twenty-ninth on the 29th. The thirtieth on the 30th. The thirty-first on the 31st.

Department of the Interior, Bureau of Land Management, Washington, D.C. 20246

[illegible]

which contribute slightly to fluctuations in the proton energy, causes a spread in the energy of the protons striking the lithium target of less than 5 kilovolts. The spread in neutron energy resulting from this resolution in proton energy relative to the spread in neutron energy caused by other reasons is very small. In fact, it is probably smaller than the probable error in the determination of the thickness of the lithium target. Therefore, its effect has been neglected.

Because of the low cross section for the formation of the metastable state, it was necessary to use a thick lithium target and to place the sample close to the target, in order to excite a significant amount of activity in the sample. This procedure involves a sacrifice in the resolution of the energy of the neutrons traversing the sample.

A target with a thickness equivalent to 60 kilovolts of proton energy was used to obtain the excitation curve shown in Figure 5. The method of determining the target thickness is described in Chapter III. The neutrons produced in a target of finite thickness follow a rectangular number-energy distribution. The average spread in neutron energy resulting from the 60-kilovolt target (expressed in proton energy) is 50 kilovolts.

The 3-cm. cadmium disks were located 4 cm. from the lithium target during the irradiations. With the sample at this position, all the neutrons emerging from the target between the angles of 0 and 20.6 degrees strike it. The spread in neutron energy, due to emission angle,

varies from 25 kilovolts at a neutron energy of 0.4 Mev to 40 kilovolts at 1.4 Mev.

In Figures 8 and 11, the activity induced in the samples is plotted against the maximum energy of the neutrons producing it. It is desirable to correct these curves so that the induced activity is plotted against the mean energy of the neutrons producing it. Willard's (W6) curves of neutron yield from the $\text{Li}^7(p,n)$ reaction versus angle of neutron emission indicate that the yield is nearly constant for angles up to 20 degrees in the energy range of this investigation. Therefore, it was assumed that the neutron flux was uniform over the whole angle at the target intercepted by the sample when making this correction.

In the following discussion, the total neutron energy spread due to target thickness is represented by ΔE_p and that due to the solid angle intercepted by the sample by ΔE_θ . The actual values of ΔE_p and ΔE_θ , for the energy range covered in this experiment, were calculated with the aid of a table, "Neutron Energy as a Function of Proton Energy and Angle for the $\text{Li}^7(p,n)\text{Be}^7$ Reaction," prepared by Willard (W6).

Let it be assumed that the cross section for the formation of the metastable state is constant over the energy spread, ΔE_θ , and that the neutron beam is monoenergetic at zero degrees ($\Delta E_p = 0$). Then, as the angle at the target increases, the increment of annular area on

where Δ is the energy of a neutron group at Δ and Δ_0 is the energy of a neutron group at Δ_0 .

In Figure 2 and 3, the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the activity is shown in the range of the activity is shown in the range of

the sample, associated with a specific increment of ΔE_0 , also increases. Therefore, the number of neutrons of a specific energy striking the sample increases as the angle at the target increases and the neutron energy decreases. The mean neutron energy, considering energy dependence on angle only, is rigorously calculated to be $[E_{\max} - 0.71 \Delta E_0]$. The scintillation counter, because of geometry, counts the decays occurring near the outside of the sample less efficiently than those occurring at the center of the sample. This tends to counteract the effect described above, and the mean energy of the effective neutrons may be taken as approximately:

$$[E_{\max} - 0.5 \Delta E_0] .$$

Then the total correction to be subtracted from E_{\max} , assuming a rectangular number-energy distribution for the spread of neutron energy due to target thickness, is approximately:

$[0.5 \Delta E_p + 0.5 \Delta E_0]$. McCue (N3) has derived a formula which includes secondary effects due to changing curvature for making this type of correction. The simplification of McCue's formula, shown below, gives an approximate, but convenient, method of making the desired correction to the neutron excitation curve.

$$A'(E_{\text{mean}}) = 4/3 A(E_{\max}) - 1/6 A(E_{\max} + \delta) - 1/6 A(E_{\max} - \delta)$$

E_{\max} = maximum neutron energy read from Figure 11.

$$\delta = 0.5 \Delta E_p + 0.5 \Delta E_0 .$$

$A(X)$ = induced activity for energy X read from Figure 11.

$E_{\text{mean}} = E_{\text{max}} - \delta$ = mean neutron energy on corrected excitation curve.

$A'(E_{\text{mean}})$ = induced activity for mean neutron energy on corrected excitation curve.

The use of the above approximation is justified because it involves only a slight deviation from the true correction, which is a small per cent of the total induced activity. This correction does not apply at the thresholds for the formation of the metastable state and higher excited levels (indicated by sharp changes in the slope of the excitation curve) or in the energy interval equal to $[\Delta E_p + \Delta E_0]$ above these thresholds. When the maximum neutron energy is just equal to the threshold energy, the correction must be zero because only neutrons with the threshold energy are contributing to the excitation. At an energy $[\Delta E_p + \Delta E_0]$ above the threshold, all the neutrons striking the sample are potentially effective, and the correction becomes applicable. In the region where the correction, according to the above formula, is not applicable, the true correction is very complicated and depends on the fraction of all the neutrons striking the sample which is effective in causing excitation, as well as on ΔE_p and ΔE_0 . The amount of experimental data obtained in these critical regions was insufficient to allow making an accurate correction to these portions of the uncorrected excitation curve of Figure 11.

4117 • *Journal of Neurosci.*, September 24, 2008 • 28(39):9911–9920

[illegible]

0750 603704

(All rights reserved.)

The use of the above procedure in testing for a difference between two groups is illustrated by the following example.

Immunization with a live attenuated virus is a

and governor will probably leave the off to the the the

and again as the economy and the location of the corporate plants

It starts off at approximately 100,000 ft above the surface and

$\left[\begin{smallmatrix} \Delta & \Delta \end{smallmatrix} \right]$ of some bounded system with all its lower subsystems with

Figure 1. (a) Gross carbon fixation rate and (b) gross carbon fixation rate per unit of leaf area.

At the University of Michigan, the researchers have been able to

Indirizzo di pubblicazione per corrispondenza: *Journal of Management Studies*, 10, 100 Brook Hill Drive, West Nyack, NY 10994, USA.

It is well known that the following identity holds:

$$\int_{\mathbb{R}^n} \Delta u \, dx = 0$$

for any function u satisfying the above conditions. This identity is a consequence of the divergence theorem and the fact that the Laplacian of a function is the divergence of its gradient.

studies on small and potentially effective, but less successful, firms.

...and ...

These results are consistent with the hypothesis that the two variables are very similar.

add additional services and file to website and on cloud for later.

Δ at all times as interface volume of particles of stable phase.

Condition (iii) is satisfied with $\delta = 1$ and $\epsilon = 1$.

...and the

provision of the amended Copyright Act of 1976.

McQue's simplified formula was applied to the excitation curve of Figure 11, except in the critical regions above the three thresholds. In these regions, the correction varies from zero at the threshold to the magnitude given by the formula at an energy of $(\Delta E_p + \Delta E_\theta)$ above the threshold. The corrected curve is shown in Figure 12. The shape of the curve in the uncertain regions above the thresholds has been estimated by extrapolating the known portions of the curve back to the threshold energies. From the cross-section standpoint, the most important part of the excitation curve is between the threshold for the formation of the metastable state and the first excited level above it. Since this portion of the curve is relatively flat, a slight error in the shape of the curve just above the threshold is not very serious. It should be noted that the threshold and the excited levels occur at the same energies on both Figure 11 and Figure 12.

The corrected excitation curve indicates that the threshold for exciting the metastable state is 400 ± 25 Kev. The presence of excited levels at 720 ± 20 Kev and at 1.15 ± 0.02 Mev is also evident. The errors assigned above are due to uncertainty in determining the target thickness, in correcting for interfering half-lives, target thickness, etc., and in drawing the curve itself.

Determination of the Absolute Cross Section for Formation of the Metastable State

A knowledge of the absolute cross section for the formation of the metastable state may facilitate assignment of a value to ℓ .




Figure 12

EXCITATION CURVE FOR 48.6 METASTABLE $^{113}\text{Cd}^{\text{III}}$ CORRECTED FOR
FINITE TARGET THICKNESS AND ANGULARITY OF INCIDENT NEUTRONS

the orbital angular momentum of the neutron which excited the state. The determination of this cross section from the experimental data already obtained requires a knowledge of the absolute neutron flux incident on the sample and of the efficiency of the scintillation counter for counting the decay of the metastable state.

The absolute neutron flux was determined for one point on the excitation curve by utilizing Hanson, Taschek, and Williams' (H7) data on neutron yield from the $\text{Li}^7(p,n)$ reaction. A sample was irradiated for one hour with neutrons from a 3.1-Mev proton beam incident on a 30-kilovolt lithium target. Hanson et al. report that a 3.1-Mev proton beam on a 40-kilovolt target yields 4.3×10^6 neutrons per microcoulomb per unit solid angle at zero degrees. Assuming that the yield is directly proportional to target thickness at this energy, the yield during the one-hour irradiation of the cadmium sample was 3.2×10^6 neutrons per microcoulomb per unit solid angle at zero degrees. The total number of microcoulombs recorded by the beam current integrator of the Rockefeller generator during this run was 20630. The solid angle intercepted by the 3-cm. diameter disk located 4 cm. away is 0.396 steradians. Therefore, 2.61×10^{10} neutrons traversed the sample during the irradiation. This sample was also counted in the scintillation counter for 45 minutes, starting 15 minutes after the end of the irradiation. The induced activity for this run, expressed as scintillation-counter counts per incident neutron was 7.56×10^{-7} . The induced activity from the excitation curve of

Figure 12 for the same mean neutron energy, 1.357 Mev, is 8.48×10^{-4} scintillation-counter counts per long-counter count. The curve of Figure 12 may be normalized to give induced activity in terms of scintillation-counter counts per incident neutron by multiplying its ordinates by the ratio $(7.56 \times 10^{-7} / 8.48 \times 10^{-4}) = 8.92 \times 10^{-4}$.

The resolving time of the scintillation counter was not short enough to discriminate between the decay of the 48.6-minute metastable state and the decay of the 8×10^{-5} -second metastable state. Therefore, each transition from the 0.396-Mev level to the 0.247-Mev level is accompanied by a simultaneous transition from the 0.247-Mev level to the ground state. Both gamma-ray transitions are partially internally converted. All the possible radiations available to the counter from this decay scheme are listed below:

- (a) 0.149-Mev unconverted gamma ray.
- (b) 0.123-Mev conversion electron.
- (c) 0.247-Mev unconverted gamma ray.
- (d) 0.221-Mev conversion electron.
- (e) 0.026-Mev cadmium X-ray.

The 0.149-Mev gamma ray is 92.25 per cent internally converted ($\alpha = 12$) and the 0.247-Mev gamma ray is 9.09 per cent internally converted ($\alpha = 0.1$) (21). The range of the 0.221-Mev electrons in cadmium is 0.005 cm. and the sample disks were 0.1625 cm. thick. Therefore, only the 0.005 cm. layer on the outside of the disk toward the counter is effective in emitting electrons, and the number from

this source is negligible. The 0.123-Mev electrons have a much shorter range, so their effect is even more negligible. The self absorption of the X-rays, 0.149-Mev gamma rays, and 0.247-Mev gamma rays in the cadmium disk is 94 per cent, 40 per cent, and 20 per cent, respectively. A comprehensive treatment of all the nuclear data given above indicates that 86 per cent of all the radiation emitted during the decay of metastable Cd^{111} is 0.247-Mev unconverted gamma rays. Since the counter efficiency probably increases with increasing gamma-ray energy, the 0.247-Mev gamma ray may be an even greater fraction of the total effective radiation. Therefore, in determining the efficiency of the scintillation counter, it will be assumed that only the 0.247-Mev gamma rays are counted. Of course, the total number of counts recorded by the scintillation counter includes 1.5-Mev beta rays from the decay of 2.6-hour Cd^{117} and 0.6-Mev beta rays, 1.13-Mev beta rays and 0.52-Mev gamma rays from the decay of 2.33-day Cd^{115} , as well as the radiation from the decay of metastable $^*\text{Cd}^{111}$. Part of the unwanted radiation from Cd^{115} and Cd^{117} is excluded by the window of the differential discriminator and the remainder has been separated from the radiation emitted by decaying metastable $^*\text{Cd}^{111}$ by the half-life-separation method already described in a previous section.

Hg^{203} decays with a 48-day half-life by emitting a 0.210-Mev beta ray and a 0.279-Mev gamma ray. This isotope was readily available at the time of this investigation. Since it emits a

[illegible]

single gamma ray of nearly the same energy as the 0.247-Mev gamma ray from metastable ^{111}Cd and also has a relatively long lifetime, Hg^{203} was chosen as a standard for estimating the efficiency of the scintillation counter for detecting the decay of metastable ^{111}Cd . Some Hg^{203} was sealed between two lucite disks, 3 cm. in diameter and 1/16 inch thick. The uniform distribution of the Hg^{203} between the disks gave approximately the same counting geometry as that of the cadmium samples. The lucite completely absorbed the beta rays. The method of determining the absolute rate of gamma-ray emission by the standard is described in Appendix B.

At the time the efficiency of the scintillation counter was determined, the Hg^{203} standard emitted 1.107×10^5 gamma rays per second. The activity of the standard, the minimum acceptable to the platinum-screen gamma-ray counter used for standardisation, was sufficient to jam the register of the differential discriminator. Therefore, an Atomic Instrument Company Model 105 decade scaler with a conventional discriminator was used with the scintillation counter for the efficiency determination. The Hg^{203} standard emitted 2.98×10^5 gamma rays during a 45-minute counting period, and the decade scaler actually recorded 1.672×10^6 counts during this time. This gives a counter efficiency of 0.56 per cent. Since the differential discriminator was used for the experimental runs, this figure must be corrected to take into account the radiation which is excluded by its window, and a correction must also be made for the radiation which is self-absorbed in the cadmium sample.

[illegible]

- (b) Determine the absolute number of decays for this specific run by applying the efficiency obtained with the decade scaler and the Hg^{203} standard.
- (c) Enter the curve of Figure 12 at the mean neutron energy used in (a) and normalize the curve to the absolute value determined in (b).

The above procedure was not used because the window of the differential discriminator excludes a large part of the unwanted activity from the decay of Cd^{115} and Cd^{117} , which is counted by the decade scaler.

The cross section for formation of the metastable state was computed by use of the following formula:

$$\frac{I_0 - I}{I_0} = 1 - e^{-n\sigma t}$$

I_0 = neutrons incident on the sample.

I = neutrons which traverse the sample without exciting the metastable state.

$I_0 - I$ = neutrons exciting the metastable state.

n = the number of scattering nuclei per cm^3 of sample.

σ = cross section for formation of the metastable state.

t = thickness of the sample.

The appropriate calculation indicates that 0.383 of all the Cd^{111} nuclei excited to the metastable level during the irradiation

The following table gives the results of the calculations for the various cases considered. The first column gives the value of α , the second column the value of β , the third column the value of γ , the fourth column the value of δ , the fifth column the value of ϵ , the sixth column the value of ζ , the seventh column the value of η , the eighth column the value of θ , the ninth column the value of ι , the tenth column the value of κ , the eleventh column the value of λ , the twelfth column the value of μ , the thirteenth column the value of ν , the fourteenth column the value of ξ , the fifteenth column the value of \omicron , the sixteenth column the value of π , the seventeenth column the value of ρ , the eighteenth column the value of σ , the nineteenth column the value of τ , the twentieth column the value of υ , the twenty-first column the value of ϕ , the twenty-second column the value of χ , the twenty-third column the value of ψ , the twenty-fourth column the value of ω , the twenty-fifth column the value of δ , the twenty-sixth column the value of ϵ , the twenty-seventh column the value of ζ , the twenty-eighth column the value of η , the twenty-ninth column the value of θ , the thirtieth column the value of ι , the thirty-first column the value of κ , the thirty-second column the value of λ , the thirty-third column the value of μ , the thirty-fourth column the value of ν , the thirty-fifth column the value of ξ , the thirty-sixth column the value of \omicron , the thirty-seventh column the value of π , the thirty-eighth column the value of ρ , the thirty-ninth column the value of σ , the fortieth column the value of τ , the forty-first column the value of υ , the forty-second column the value of ϕ , the forty-third column the value of χ , the forty-fourth column the value of ψ , the forty-fifth column the value of ω , the forty-sixth column the value of δ , the forty-seventh column the value of ϵ , the forty-eighth column the value of ζ , the forty-ninth column the value of η , the fiftieth column the value of θ , the fifty-first column the value of ι , the fifty-second column the value of κ , the fifty-third column the value of λ , the fifty-fourth column the value of μ , the fifty-fifth column the value of ν , the fifty-sixth column the value of ξ , the fifty-seventh column the value of \omicron , the fifty-eighth column the value of π , the fifty-ninth column the value of ρ , the sixtieth column the value of σ , the sixty-first column the value of τ , the sixty-second column the value of υ , the sixty-third column the value of ϕ , the sixty-fourth column the value of χ , the sixty-fifth column the value of ψ , the sixty-sixth column the value of ω , the sixty-seventh column the value of δ , the sixty-eighth column the value of ϵ , the sixty-ninth column the value of ζ , the seventieth column the value of η , the seventy-first column the value of θ , the seventy-second column the value of ι , the seventy-third column the value of κ , the seventy-fourth column the value of λ , the seventy-fifth column the value of μ , the seventy-sixth column the value of ν , the seventy-seventh column the value of ξ , the seventy-eighth column the value of \omicron , the seventy-ninth column the value of π , the eightieth column the value of ρ , the eighty-first column the value of σ , the eighty-second column the value of τ , the eighty-third column the value of υ , the eighty-fourth column the value of ϕ , the eighty-fifth column the value of χ , the eighty-sixth column the value of ψ , the eighty-seventh column the value of ω , the eighty-eighth column the value of δ , the eighty-ninth column the value of ϵ , the ninetieth column the value of ζ , the ninety-first column the value of η , the ninety-second column the value of θ , the ninety-third column the value of ι , the ninety-fourth column the value of κ , the ninety-fifth column the value of λ , the ninety-sixth column the value of μ , the ninety-seventh column the value of ν , the ninety-eighth column the value of ξ , the ninety-ninth column the value of \omicron , the hundredth column the value of π .

decay during the 45-minute counting period. Therefore, the quantity $(I_0 - I)$ also equals the absolute number of decays occurring during the counting period divided by 0.383. After final simplification:

$$\frac{I_0 - I}{I_0} = 0.656 \times \text{ordinate of Figure 12.}$$

Figure 13 is a graph of the cross section for formation of the metastable state plotted against the mean energy of the incident neutrons. The ordinate of this curve is subject to an estimated error of ± 25 per cent due to:

- (a) The probable error in the ordinate of Figure 12.
- (b) The uncertainty in the absolute rate of gamma-ray emission by the Hg^{203} standard.
- (c) The assumption that the scintillation counter counts only the 0.247-Mev gamma ray from the decay of the metastable state in Cd^{111} .
- (d) Error in determining the absolute neutron flux.

Comparison of the Theoretical and Experimental Cross Sections

A comparison of the experimental cross section for the formation of the metastable state with the theoretical cross section, $\sigma_0(l)$, for the formation of the compound nucleus, Cd^{112} , as a function of l for the incoming neutron allows the assignment of a maximum value to the orbital angular momentum the neutron can carry in. The quantity,

the reaction was stopped by 5% NaOH solution and the mixture was extracted with 10 ml of diethyl ether. The ether was dried over anhydrous CaCl_2 and the solvent was removed by distillation. The residue was purified by distillation under reduced pressure. Yield: 1.5 g (10%).

$$M_{\text{total}} = M_{\text{solid}} + M_{\text{liquid}} = \frac{1}{2} \rho V$$


To collect the money from the sale of the property, the following is the list of the property to be sold:

- (a) The results show in the column of Page 11.
- (b) The results in the column of Page 11.
- (c) The results in the column of Page 11.
- (d) The results in the column of Page 11.
- (e) The results in the column of Page 11.
- (f) The results in the column of Page 11.
- (g) The results in the column of Page 11.
- (h) The results in the column of Page 11.
- (i) The results in the column of Page 11.
- (j) The results in the column of Page 11.
- (k) The results in the column of Page 11.
- (l) The results in the column of Page 11.
- (m) The results in the column of Page 11.
- (n) The results in the column of Page 11.
- (o) The results in the column of Page 11.
- (p) The results in the column of Page 11.
- (q) The results in the column of Page 11.
- (r) The results in the column of Page 11.
- (s) The results in the column of Page 11.
- (t) The results in the column of Page 11.
- (u) The results in the column of Page 11.
- (v) The results in the column of Page 11.
- (w) The results in the column of Page 11.
- (x) The results in the column of Page 11.
- (y) The results in the column of Page 11.
- (z) The results in the column of Page 11.

Copyright © 2004 by John Wiley & Sons, Inc.

of the engine which is the most common type of engine used in the world. The engine is a four stroke engine and is of the type which is used in the most common type of engine used in the world. The engine is a four stroke engine and is of the type which is used in the most common type of engine used in the world.

Figure 13
CROSS SECTION FOR FORMATION OF
METASTABLE STATE IN Cd^{III}




$\sigma_0(l)$, has been calculated following the method of Blatt and Weisskopf (23), as described in Chapter II, for the neutron-energy region from 0.2 Mev to 1.2 Mev for values of l from zero through 5. It should be noted that these calculations involve the kinetic energy of the neutron in the center of mass system. However, since Cd^{111} is a relatively heavy nucleus, the kinetic energy of the neutron is essentially the same in both the center of mass and the laboratory system. These $\sigma_0(l)$'s which are in a range a few orders of magnitude above and below the experimental cross section for the formation of the metastable state are plotted against mean neutron energy in Figure 14. The experimental cross section in the vicinity of the threshold has also been plotted on the graph in Figure 14. The curves of Figure 14 are plotted on semi-log graph paper in order that a practical scale may be used for the vertical coordinate. An unqualified evaluation of these curves indicates that the most angular momentum which a neutron can carry into the compound nucleus is $l = 3$.

The formula for W_1 , the fraction of nuclei decaying from the compound nucleus to a specific excited state, is also given in Chapter II. An inspection of the formulas for $\sigma_0(l)$ and W_1 shows that for the case of $l = 0$, the cross section $\sigma_0(0)$ decreases as $(E - E_{th})^{-1/2}$, and, therefore, the probability of decay to the specific state associated with W_1 increases as $(E - E_{th})^{1/2}$. When an $l = 1$ neutron is emitted in the decay of the compound nucleus, the probability of

$\sigma(l)$, has been estimated following the method of Blatt and
 Blatt (1952) as described in Chapter II, for the neutron-energy
 region from 0.5 meV to 1.5 meV the value of l is
 2. It should be noted that these calculations involve the kinetic
 energy of the neutron in the center of mass system. However, since
 the l is a relatively heavy nucleus, the kinetic energy of the
 neutron is essentially the same in both the center of mass and the
 laboratory system. Thus $\sigma(l)$, which was in a range a few orders
 of magnitude above and below the experimental cross section for the
 formation of the metastable state are plotted against mean neutron
 energy in Figure 10. The experimental cross section in the vicinity
 of the threshold has also been plotted on the graph in Figure 10.
 The curves of Figure 10 are plotted on semi-log paper in order
 that a horizontal scale may be used for the vertical coordinate. It
 is pointed out that the experimental cross section is not a
 constant which is constant and may have the experimental value is
 $l = 2$

The formula for l , the fraction of nuclei decaying from
 the compound nucleus in a specific excited state, is also given in
 Chapter II. In the case of the formula for $\sigma(l)$ and σ above
 that for the case of $l = 0$, the cross section $\sigma(0)$ becomes
 $(1 - \frac{1}{2} \frac{1}{l})^{1/2}$ and therefore the probability of decay to the ground
 state becomes $1 - \frac{1}{2} \frac{1}{l}$ as $l \rightarrow \infty$.

It is noted in the case of the compound nucleus, the probability of



decay increases as $(E - E_{th})^{3/2}$ and, in general, as $(E - E_{th})^{(l+1/2)}$.

A mathematical analysis of these formulas indicates that if the decay occurs by the emission of an $l = 0$ neutron, the curve of W_1 versus neutron energy approaches its threshold perpendicular to the horizontal axis. For the case of higher values, the curve approaches the threshold tangent to the horizontal axis.

When the experimental cross section for the formation of the metastable state is divided by the cross section for the formation of the compound nucleus, the resulting quantity is W_1 , the probability that the compound nucleus will decay to the metastable state. Such a calculation was carried out using the experimental cross section for the formation of the metastable state and the theoretical value of $\sigma_0(3)$. The curve of W_1 versus mean neutron energy is shown in Figure 15. The shape of this curve in the vicinity of the threshold indicates that an $l = 0$ neutron is emitted from the compound nucleus. The curve deviates from the $(E - E_{th})^{1/2}$ curve as the neutron energy increases above the threshold value, because competition from decay to levels other than the metastable state commences as the energy increases.

Mean Neutron Energy - Mev

Figure 15




Figure 15

V. CONCLUSIONS AND SUGGESTIONS FOR FURTHER INVESTIGATION

DISCUSSION AND SUMMARY OF RESULTS

The neutron excitation curve obtained during this investigation indicates that the threshold energy for the formation of the metastable state is 0.400 Mev and also reveals the existence of energy levels at 0.720 Mev and 1.150 Mev. The location of the metastable state is currently accepted as 0.396 Mev above the ground state. Therefore, the neutron excitation curve indicates the direct excitation of the metastable state. The existence of the 0.720-Mev energy level has not been reported previously. The 1.150-Mev energy level is probably the same one which Wiedenbeck (W7) obtained by both photon and electron bombardment and to which he assigned the energy of 1.250 Mev. The results of this investigation and those results obtained by Ebel (E1) indicate that neutron excitation gives slightly lower values of energy than photon or electron excitation gives for what are apparently the same levels.

The comparison of theoretical and experimental cross sections indicates that an $\ell = 3$ neutron forms the compound nucleus, which then emits an $\ell = 0$ neutron. If the metastable state is excited directly, conservation of angular momentum and parity for the above ℓ values gives the metastable state a possible angular momentum of $9/2$, $7/2$, or $5/2$ and a parity different than that of the ground state. According to Figure 5, the most probable value of

angular momentum for the metastable state, when its parity is different than that of the ground state, is $13/2$ and the 0.149-Mev transition is by electric 2^4 -pole radiation, while the 0.247-Mev transition is by magnetic quadrupole radiation. An angular momentum of $9/2$, $7/2$, or $5/2$ for the metastable state is possible for this decay scheme, but much less probable than $13/2$. The probability for the above possible values decreases rapidly as the magnitude of the angular momentum decreases.

Although the curve for the experimental cross section shown in Figure 14 is above the theoretical curve for $l = 4$, it is much closer to the curve for $l = 4$ than the curve for $l = 3$. After considering the uncertainties involved in determining the experimental cross section, it is conceivable that the experimental curve may be so much in error that its true position should be below the $l = 4$ theoretical curve. If this is the case, an $l = 4$ neutron excites the metastable state directly. If the points on the experimental cross section curve retain their relative positions and the curve, as a whole, is moved vertically downward until it is below the $l = 4$ theoretical curve, a new graph similar to the one in Figure 15 would still indicate that an $l = 0$ neutron is emitted by the compound nucleus. These new conditions would give the metastable state a possible angular momentum of $11/2$, $9/2$, or $7/2$ and the same parity as the ground state. An angular momentum of $11/2$ and no parity change are consistent and most probable, as shown in Figure 5, for the assignment of electric 2^4 -pole radiation to the 0.149-Mev transition

exactly known for the magnetic state when the ratio is either
 not less than of the critical value, is $1/2$ and the 0.100- μ transition
 is in the h -pole position, while the 0.200- μ transition is in
 magnetic h -pole position. An angular momentum of $0/2$, $1/2$, or $3/2$
 for the magnetic state is possible for both cases, but more
 likely probably than $1/2$. The possibility for the other possible
 values depends mainly on the nature of the angular momentum
 quantum.

Although the curve for the experimental cross section shows
 in Figure 10 to have the maximum curve for $l = 4$, it is more
 likely in the curve for $l = 4$ than the curve for $l = 3$. After con-
 sidering the uncertainties involved in determining the experimental
 cross section, it is probable that the experimental curve can be
 well fitted by a curve with $l = 4$ and the curve for $l = 3$
 is also a good fit. It is in the same way, as l increases, the
 fit becomes even better. If the points on the experimental
 cross section curve were their relative positions and the curve
 as a whole, it would probably decrease until it is below the $l = 4$
 theoretical curve, a very good fit to the one in Figure 10 would
 still indicate that $l = 0$ would be ruled by the experiment.
 However, these two positions would give the magnetic state a pos-
 sible angular momentum of $1/2$, $3/2$, or $5/2$ and the same parity as
 the critical state. An angular momentum of $1/2$ and an odd parity
 are excluded and most probable, as shown in Figure 9, for the
 assignment of electric h -pole radiation to the 0.100- μ transition.

and magnetic dipole radiation to the 0.247-Mev transition. The above values of angular momentum are also possible, but not probable, for the case where the 0.149-Mev transition is by electric 2^4 -pole radiation and the 0.247-Mev transition is by electric quadrupole radiation.

McGinnis (14) has very recently stated that the 0.396-Mev level has an angular momentum of $13/2$ and decays by electric 2^4 -pole radiation to the 0.247-Mev level and that the 0.247-Mev level has an angular momentum of $5/2$ and decays by electric quadrupole radiation. These assignments of spin and type of multipole give both metastable states the same parity as the ground state. His determinations are based on a comparison of experimental values of half-life, conversion ratios, etc., with theoretical relations presented in Rose's (11) tables for the K conversion coefficient. (Rose's tables were not available to the author until after the interpretation of the experimental results of this investigation had been accomplished). McGinnis's decay scheme is the one which the author initially considered most probable (see Figures), but unfortunately it does not agree with the most probable values of angular momentum indicated as possible by the experimental results of this investigation.

In^{111} decays by electron capture to a 0.420-Mev level in Cd^{111} . The probable error in the threshold energy obtained by this investigation for exciting the metastable state is of such magnitude that the excitation curve could be read to indicate the direct excitation of the 0.420-Mev level and its immediate decay to the metastable state. The presence of the 0.420-Mev level has only been observed in

[illegible][illegible][illegible]

1. The first step in the process is to identify the problem or issue that needs to be addressed. This involves gathering information and understanding the context of the situation.

connection with the decay of In^{111} . Attempts to separate any 48.6-minute metastable ^{111}Cd from decaying In^{111} have been unsuccessful (M4). McGinnis has assigned an angular momentum of $7/2$ to this level and reports that it decays to the 0.247-Mev level by the emission of magnetic dipole radiation. If the 0.420-Mev level was excited directly during this investigation by an $l = 3$ neutron, the parity considerations of McGinnis's decay scheme for this level are not satisfied. Direct excitation of the 0.420-Mev level by an $l = 4$ neutron is consistent with parity considerations, but very improbable. Since the energy differential between the 0.420-Mev level and the 0.396-Mev level is so small, a transition from the 0.420-Mev level to the 48.6-minute metastable level is even more improbable. Therefore, it is very doubtful that the 0.420-Mev level was excited directly.

Figure 16 shows the most probable decay schemes of those considered possible as a result of this investigation.

Suggestions for Further Investigation

The neutron excitation curve should be extended to higher energies in order to confirm the energy levels at 1.68, 2.08, and 2.56 Mev obtained by Wiedenbeck (W7) from both photons and electron excitation.

It is planned to retain the experimental equipment, used during this investigation, intact for some time into the future. Therefore, it is recommended that additional experimental work be directed toward a better determination of the absolute cross section for the

[illegible]

The level of the water in the reservoir is 100 feet above the level of the sea. The level of the water in the reservoir is 100 feet above the level of the sea. The level of the water in the reservoir is 100 feet above the level of the sea.




It is not possible to determine the exact date of the
the above mentioned events. The only information
available is that the events took place in the
month of May, 1944.

1945-1946

... ..




It is noted in the report that the investigation was conducted by the FBI and the results are being reported to the Department of Justice.

$$\ell_{n-in} = 3 \quad \ell_{n-out} = 0$$




	Energy	Parity	Ang Mom
	396 Mev	\mp	$9/2$
	247	\mp	$5/2$ or $3/2$
	0	\pm	$1/2$

CASE I

$$\ell_{n-in} = 4 \quad \ell_{n-out} = 0$$

	Energy	Parity	Ang Mom
	396 Mev	\pm	$11/2$
	247	\pm	$3/2$
	0	\pm	$1/2$

CASE II

	Energy	Parity	Ang Mom
	396 Mev	\pm	$11/2$
	247	\pm	$5/2$ or $3/2$
	0	\pm	$1/2$

CASE III

Figure 16

MOST PROBABLE ASSIGNMENT OF ANGULAR MOMENTUM AND PARITY TO THE METASTABLE STATE OF Cd^{111} .

formation of the metastable state. This effort should include a confirmation of the absolute neutron flux incident on the sample by a method independent of Hanson, Taschek and Williams (H7), a more accurate method of determining the efficiency of the scintillation counter, and a careful experimental analysis, by other means than half-life separation, of the different kinds of radiation detected by the counter during the decay of metastable ^{111}Cd .

If the scintillation counter is moved to a location in the near vicinity of the target room of the Rockefeller generator, an investigation of the 2.3-minute metastable ^{113}Cd , similar to the one described in this report, may be made. Because of the relatively short half-life of metastable ^{113}Cd , the time required for irradiations and counting will be much less than that spent during the investigation of metastable ^{111}Cd . The author has procured disks of barium peroxide which may be used for another similar investigation of the excitation of the 2.6-minute metastable state in Ba^{137} by fast neutron bombardment. It seems that the investigations of Cd^{113} and Ba^{137} could be conveniently carried on at the same time.

Appendix U contains a brief summary of several unsuccessful attempts to obtain neutron excitation curves for metastable states in various isomers. This work was carried on before the final selection of Cd^{111} as a subject for this investigation was made.

BIBLIOGRAPHY

- A1 P. Axel and S. Dancoff, Phys. Rev. 76, 892 (1949).
- B1 S. W. Barnes and P. W. Aradine, Phys. Rev. 55, 50 (1939).
- B2 H. Bethe, Rev. Mod. Phys. 9, 220 (1937).
- B3 J. M. Blatt and V. F. Weisskopf, Laboratory for Nuclear Science and Engineering, M.I.T., Technical Report No. 42.
- D1 S. M. Dancoff and P. Morrison, Phys. Rev. 55, 122 (1939).
- D2 M. Deutsch and D. Stevenson, Phys. Rev. 76, 154 (1949).
- D3 M. Deutsch and W. Wright, Phys. Rev. 77, 130 (1950).
- D4 A. J. Dempster, Phys. Rev. 71, 144 (1947).
- E1 A. A. Ebel, "Metastable States in Medium and Heavy Weight Nuclei," Ph. D. Thesis, M.I.T., (1950).
- E2 W. C. Elmore and M. L. Sands, Electronics: Experimental Techniques, National Nuclear Energy Series V-1, (New York: McGraw-Hill Book Co., Inc., 1949).
- G1 M. Goldhaber, R. D. Hill, and L. Szilard, Phys. Rev. 55, 47 (1939).
- G2 M. Goldhaber and G. O. Nuelhouse, Phys. Rev. 74, 1245 (1949).
- H1 H. Høle, Arkiv. Mat. Astron. Fysik, 36A, No. 9, (1948).
- H2 H. H. Hebb and G. E. Uhlenbeck, Physica 5, 605 (1938).
- H3 F. M. Hadden, Ph. D. Thesis, M.I.T. (1950).
- H4 A. Hanson and J. McKibben, Phys. Rev. 72, 673 (1947).
- H5 A. Helmholtz, R. Hayward, and C. L. McGinnis, Phys. Rev. 75, 1469 (1949).
- H6 H. H. Hebb and E. Nelson, Phys. Rev. 58, 489 (1940).
- H7 A. Hansen, R. Taschek, and J. M. Williams, Rev. Mod. Phys. 21, 635 (1949).

CONTENTS

1. *General Principles of the Theory of the* (1901)
2. *On the Theory of the* (1902)
3. *On the Theory of the* (1903)
4. *On the Theory of the* (1904)
5. *On the Theory of the* (1905)
6. *On the Theory of the* (1906)
7. *On the Theory of the* (1907)
8. *On the Theory of the* (1908)
9. *On the Theory of the* (1909)
10. *On the Theory of the* (1910)
11. *On the Theory of the* (1911)
12. *On the Theory of the* (1912)
13. *On the Theory of the* (1913)
14. *On the Theory of the* (1914)
15. *On the Theory of the* (1915)
16. *On the Theory of the* (1916)
17. *On the Theory of the* (1917)
18. *On the Theory of the* (1918)
19. *On the Theory of the* (1919)
20. *On the Theory of the* (1920)

- J1 B. Jennings, Proc. I.R.E. 38, 1126 (1950).
- L1 K. Lark-Horovitz, J. R. Risser, and R. M. Smith, Phys. Rev. 55, 878 (1939).
- L2 I. Loew, Phys. Rev. 59, 835 (1941).
- L3 J. Lovington, M.S. Thesis, M.I.T. (1951).
- M1 P. B. Moon, "Artificial Radioactivity," Cambridge University Press (1949).
- M2 B. Moyer, B. Peters, and F. H. Schmidt, Phys. Rev. 69, 666 (1946).
- M3 J.J.G. McGue, private communication.
- M4 C. L. McGinnis, Phys. Rev. 81, 734 (1951).
- N1 "Nuclear Data," NBS 499, National Bureau of Standards, U.S. Department of Commerce.
- P1 W. M. Preston and Clark Goodman, Bul. Amer. Phys. Soc. 26, 29 (1951).
- P2 W. C. Peacock, "A Study of the Absolute Efficiency of Gamma-Ray Counters with Application to Nuclear Spectroscopy," Ph. D. Thesis, M.I.T. (1944).
- R1 M. Z. Rose, G. H. Goertzel, W. I. Spinrad, J. Harr, and P. Strong, Phys. Rev. 76, 1883 (1949).
- S1 O. T. Seaborg and I. Pearlman, Rev. Mod. Phys. 20, 585 (1948).
- S2 W. H. Sullivan, Trilinear Chart of Nuclear Species, (New York: John Wiley and Sons, 1949).
- S3 E. Segre and A. O. Holmhelts, Rev. Mod. Phys. 21, 271 (1949).
- S4 W. A. Schoenfeld and R. W. Duborg, M.S. Thesis, M.I.T. (1951).
- T1 H. M. Taylor and H. F. Mott, Proc. Roy. Soc. A142, 215 (1933).
- T2 R. Taschek and A. Remmendinger, Phys. Rev. 74, 373 (1948).
- W1 M. L. Wiedenbeck, Phys. Rev. 67, 92 (1945).
- W2 M. L. Wiedenbeck, Phys. Rev. 67, 267 (1945).

1. J. L. Thompson, *Proc. I.R.S.* 22 (1930).
2. J. L. Thompson, J. A. Brown, and R. C. Smith, *Proc. Roy. Soc.* 121 (1928).
3. J. L. Thompson, *Proc. Roy. Soc.* 121 (1928).
4. J. L. Thompson, *Proc. Roy. Soc.* 121 (1928).
5. J. L. Thompson, *Proc. Roy. Soc.* 121 (1928).
6. J. L. Thompson, *Proc. Roy. Soc.* 121 (1928).
7. J. L. Thompson, *Proc. Roy. Soc.* 121 (1928).
8. J. L. Thompson, *Proc. Roy. Soc.* 121 (1928).
9. J. L. Thompson, *Proc. Roy. Soc.* 121 (1928).
10. J. L. Thompson, *Proc. Roy. Soc.* 121 (1928).
11. J. L. Thompson, *Proc. Roy. Soc.* 121 (1928).
12. J. L. Thompson, *Proc. Roy. Soc.* 121 (1928).
13. J. L. Thompson, *Proc. Roy. Soc.* 121 (1928).
14. J. L. Thompson, *Proc. Roy. Soc.* 121 (1928).
15. J. L. Thompson, *Proc. Roy. Soc.* 121 (1928).
16. J. L. Thompson, *Proc. Roy. Soc.* 121 (1928).
17. J. L. Thompson, *Proc. Roy. Soc.* 121 (1928).
18. J. L. Thompson, *Proc. Roy. Soc.* 121 (1928).
19. J. L. Thompson, *Proc. Roy. Soc.* 121 (1928).
20. J. L. Thompson, *Proc. Roy. Soc.* 121 (1928).

- W3 M. L. Wiedenbeck, Phys. Rev. 67, 53 (1945).
- W4 M. L. Wiedenbeck, Phys. Rev. 68, 1 (1945).
- W5 B. Walden and M. L. Wiedenbeck, Phys. Rev. 63, 60 (1943).
- W6 H. D. Willard, "Interaction of Fast Neutrons with Nuclei," Ph. D. Thesis, M.I.T. (1950).
- W7 M. L. Wiedenbeck, Phys. Rev. 66, 36 (1944).

1891. The first year of the century.

1892. The second year of the century.

1893. The third year of the century.

1894. The fourth year of the century.

1895. The fifth year of the century.

1896. The sixth year of the century.

1897. The seventh year of the century.

1898. The eighth year of the century.

1899. The ninth year of the century.

1900. The tenth year of the century.

1901. The eleventh year of the century.

1902. The twelfth year of the century.

1903. The thirteenth year of the century.

1904. The fourteenth year of the century.

1905. The fifteenth year of the century.

1906. The sixteenth year of the century.

1907. The seventeenth year of the century.

1908. The eighteenth year of the century.

1909. The nineteenth year of the century.

1910. The twentieth year of the century.

ACKNOWLEDGMENTS

The graduate study of the author was sponsored by the Bureau of Ships and the U.S. Naval Postgraduate School. The work reported herein was undertaken in conjunction with the Nuclear Shielding Project of the Laboratory for Nuclear Science and Engineering, M.I.T., which project is sponsored by the Bureau of Ships and the Office of Naval Research.

The author wishes to express his appreciation to Dr. Clark Goodman, supervisor of this thesis, for his advice, interest, and constructive criticisms during the progress of this research and the preparation of this thesis; to Dr. J.J.G. McQue for the unselfish contribution of a very large portion of his time to assisting with the experimental phases of this research and to making constructive criticisms during the preparation of this thesis; to Mr. Leonard Hubacher for many valuable discussions of theory; to Dr. Donald Stevenson and Mr. Marvin VanDilla for their suggestions on both the theory and the experimental procedure; to Dr. Paul Stelson, Mr. Donald Thompson, Mr. Ira Slawson, and Mr. John Adams for their aid in conducting the experiments at the Rockefeller generator; to Mrs. Evelyn McKinley for typing the manuscript; and to Mrs. Grace Rowe for reproducing the figures.

The preparation of the cadmium, rhodium, columbium, and mercury samples by Dr. T. S. Hagel, of the M.I.T. Metallurgical Project, is appreciated.

© 2000 Blackwell Science Ltd

The estimate of the value of the property was prepared by the Bureau of Internal Revenue and the Bureau of Customs. The value of the property was estimated at \$100,000.00. The value of the property was estimated at \$100,000.00.

The writer wishes to express his appreciation to Dr. Clark
Benson, president of this Society, for his advice, interest, and en-
couragement during the progress of this research and the
preparation of this thesis; to Dr. J. L. G. Baker for the excellent con-
struction of a very large portion of his data in connection with the
experimental phases of this research and in making constructive criti-
cisms during the preparation of this thesis; to Dr. Leonard B. Bowers
for many valuable discussions of theory; to Dr. Donald Stevenson and
Dr. Lewis Phillips for their suggestions on both the theory and the
experimental procedure; to Dr. Paul Wilson, Dr. Donald Thompson, Dr.
L. A. Brown, and Dr. John Adams for their aid in conducting the ex-
periments at the Rochester Community College; to Mrs. Betty Hedges for
typing the manuscript; and to Mrs. Grace Reed for reproducing the
figures.

[illegible]

APPENDIX A

ALTERATION OF THE SINGLE-CHANNEL DIFFERENTIAL DISCRIMINATOR

The basic wiring diagram for the single-channel differential discriminator is the Laboratory for Nuclear Science and Engineering Drawing No. D-741-A, File 6420. This drawing calls for a 4700-ohm resistance in the plate circuit of the first tube of the discriminator which sets the low side of the window (6A07). This resistance was reduced to 3700 ohms by placing an 1000-ohm resistance in parallel with the original 4700 ohms. This alteration reduced the hysteresis in the discriminator circuit; i.e., the signal voltage at which the discriminator recovered after being triggered was reduced. Such a change lowered the minimum setting at which the discriminator would operate.


APPENDIX B

DETERMINATION OF THE ABSOLUTE RATE OF GAMMA-RAY EMISSION
OF THE Hg^{203} STANDARD

A platinum-screen gamma-ray counter, belonging to the Radioactivity Center of the M.I.T. Physics Department, was used for estimating the absolute rate of gamma-ray emission by a sample of Hg^{203} . This sample was eventually used in estimating the efficiency of the scintillation counter for detecting the decay of metastable ^{111}Cd . The counter is similar to the platinum-screen counter investigated by Pencoek (P2) during his determination of gamma-ray counter efficiencies.

A table of factors has been compiled by the Radioactivity Center for use of the platinum-screen counter with a number of different isotopes. This table makes it possible to determine the absolute number of disintegrations per unit time in a sample of any one of the isotopes by comparison of its counting rate with that of a radium standard when both are located the same distance from the counter. The factors for several of the isotopes with relatively simple gamma-ray decay schemes are plotted against the total energy of the gamma rays emitted in Figure 17. The factor for Hg^{203} was obtained by extrapolating this curve. The Hg^{203} sample and a radium standard were counted for equal periods of time at equal distances from the counter. Runs were made at several distances from the counter to insure that

Figure 17
 CALIBRATION CURVE FOR STANDARDIZING
 Hg^{203} SAMPLE



the geometry of the sample and the radium standard was approximately the same. The instructions for using the factor are: Compare the net counting rates from the sample and the radium standard at the same distance from the gamma-ray counter. The net counting rate for the sample, multiplied by the factor, is to the net counting rate for the standard as the ratio of their absolute disintegration rates. The extrapolation process is subject to considerable error, and the use of an isotope as a standard for which the factor was definitely known would have been more desirable. However, none of the calibrated isotopes had total gamma-ray energies approximating the single 0.247-Mev gamma ray predicted as comprising 86 per cent of the decay radiation from metastable ^{111}Cd .

[illegible]

APPENDIX C

UNSUCCESSFUL PRELIMINARY EXPERIMENTS IN PRODUCING AND DETECTING
THE DECAY OF METASTABLE STATES

Samples of Rh^{103} were irradiated with fast neutrons for one hour, but the decay of the 53-minute metastable state was not detected. This isotope is 100 per cent abundant, and from the per cent abundance standpoint, it should be an ideal subject for this type of investigation because no interfering decays will occur. The metastable state decays by the emission of a 0.032-Mev gamma ray. Evidently the cross section for the formation of the metastable state is very small or the counting equipment was not sensitive enough to detect the low energy radiation emitted. Wiedenbeck (42) reports observing the decay of metastable ^{103}Rh with delicate counters made of rhodium.

Samples of Os^{93} were irradiated with fast neutrons for one hour, but the decay of the 42-day metastable state was not detected. This failure was probably due to the relatively short time of irradiation. Use of the Rockefeller generator for a period of time comparable to the half-life of the metastable state is obviously impossible.

A sample of mercury deposited on a gold backing was irradiated with fast neutrons for one hour, but decay of the 43.6-minute metastable state in Hg^{201} was not detected. The failure was probably due to a small cross section for the formation of the metastable state and an insufficient abundance of the isotope, Hg^{201} , in natural mercury.

APPENDIX B

ANALYSIS OF THE DATA OBTAINED FROM THE INVESTIGATION

THE DATA ON THE INVESTIGATION

The data obtained from the investigation are as follows:

1. The data on the investigation are as follows:

2. The data on the investigation are as follows:

3. The data on the investigation are as follows:

4. The data on the investigation are as follows:

5. The data on the investigation are as follows:

6. The data on the investigation are as follows:

7. The data on the investigation are as follows:

8. The data on the investigation are as follows:

9. The data on the investigation are as follows:

10. The data on the investigation are as follows:

11. The data on the investigation are as follows:

12. The data on the investigation are as follows:

13. The data on the investigation are as follows:

14. The data on the investigation are as follows:

15. The data on the investigation are as follows:

16. The data on the investigation are as follows:

17. The data on the investigation are as follows:

18. The data on the investigation are as follows:

19. The data on the investigation are as follows:

JUL 2
FEB 5

BINDERY
155

15626

Thesis
F73

Francis


Isomeric metastable states in
medium and heavy weight nuclei.

Library
U. S. Naval Postgraduate School
Monterey, California



thesF73

Isomeric metastable states in medium and



3 2768 001 95977 8

DUDLEY KNOX LIBRARY